

The Yin and Yang of Cannabis-induced Psychosis: the Actions of Δ^9 -Tetrahydrocannabinol and Cannabidiol in Rodent Models of Schizophrenia

J.C. Arnold^{1,2,3,*}, A.A. Boucher² and T. Karl^{3,4,5}

¹Discipline of Pharmacology, University of Sydney NSW 2006, Australia; ²Brain and Mind Research Institute, Sydney, NSW 2050, Australia; ³Schizophrenia Research Institute, Darlinghurst NSW 2010, Australia; ⁴Neuroscience Research Australia, Randwick, NSW 2031, Australia; ⁵University of New South Wales, Sydney, NSW 2052, Australia

Abstract: The link between cannabis and psychosis has often been debated with polarized views on the topic. There is substantial epidemiological evidence showing that cannabis increases the risk of psychosis, whereas other research suggests that schizophrenia patients self-medicate with the substance. These conflicting accounts may at least be partially explained by the two phytocannabinoids cannabidiol (CBD) and Δ^9 -tetrahydrocannabinol (THC) and their opposing actions on schizophrenia-related symptoms. In the present review we will first focus on how traditional rodent models of schizophrenia have been used to improve our understanding of the pro-psychotic actions of THC and the anti-psychotic actions of CBD. We will also review novel rodent models used to address genetic vulnerability to cannabis-induced schizophrenia and show that specific genes are being uncovered that modulate cannabinoid action (e.g. the schizophrenia susceptibility gene neuregulin 1). We will also review rodent studies that have addressed interactions between THC and CBD. These animal studies underscore great complexity with some studies showing that CBD antagonises the neurobehavioural effects of THC, while others show the opposite, that CBD potentiates the actions of THC. Various mechanisms are put forth to explain these divergent effects such as CBD antagonism at central CB1 receptors or that CBD inhibits proteins that regulate THC disposition and metabolism (e.g. the ABC transporter, P-glycoprotein).

Keywords: Δ^9 -tetrahydrocannabinol, cannabidiol, schizophrenia, rodent model, CB1, CB2, neuregulin.

INTRODUCTION

It has been estimated that psychotic disorders affect 3% of the world's population [1]. Those diagnosed specifically with schizophrenia approaches 1%, although there exist a range of related diagnoses such as schizophreniform disorder, schizoaffective disorder, delusional disorder and drug-induced psychosis that can also be added to estimates of lifetime prevalence of psychosis. The onset of schizophrenia occurs normally in late adolescence/early adulthood and is characterized by negative symptoms (e.g. social withdrawal), cognitive dysfunction and positive symptoms (e.g. hallucinations and delusions). The chronic nature of schizophrenia makes it one of the leading forms of permanent disability. Since the discovery of typical antipsychotic drugs, the subsequent development of atypical compounds has provided marginal improvement in the treatment of schizophrenia. These agents are variably effective in dampening positive symptoms and 30% of patients remain treatment-resistant [2]. Importantly, antipsychotic drugs are poorly effective in treating cognitive dysfunction and negative symptoms in schizophrenia [3].

Although the exact causes of schizophrenia are still unknown, the role played by cannabis as a major risk factor has been widely debated with polarized views on the topic. For example, there is substantial epidemiological evidence showing that cannabis increases the risk of psychosis by at least 2-fold, whereas other research suggests that schizophrenia patients self-medicate with the substance to reduce schizophrenia symptoms or to alleviate the side effects of antipsychotic medication. These conflicting accounts may be partially explained by two phytocannabinoids cannabidiol (CBD) and Δ^9 -tetrahydrocannabinol (THC) and their opposing actions on schizophrenia-related symptoms. Indeed much human and animal research is now focussing on these two constituents (see also other papers published in this issue). A growing body of evidence indicates that THC increases the risk of anxiety, psychotic symptoms and memory impairment in healthy individuals

[4, 5] and those with an established psychotic disorder such as schizophrenia [6, 7]. In contrast, CBD, another major constituent of cannabis that lacks detectable intrinsic psychoactivity, has anxiolytic [8-10] and possibly antipsychotic properties [11, 12] and does not appear to impair memory or other cognitive functions. Furthermore, when co-administered, CBD can reduce the anxiogenic, memory-impairing and psychotomimetic effects induced by THC [13].

Other papers in the present issue will specifically address the neurobiological effects of cannabinoids on the brain. Functional neuroimaging studies directly comparing THC and CBD found distinct modulatory effects on regional neural responses to fearful faces [14]. Specifically, the authors observed a CBD-induced attenuation of neural responses to intensely fearful faces in the amygdala and cingulate cortex, which was correlated with an electrophysiological response and behavioural evidence for an anxiolytic effect. There was also a distinct effect for CBD on the brain connectivity linking these two regions [15]. In a subsequent fMRI study in the same cohort, THC and CBD had opposing effects on striatal activation during verbal recall, on hippocampal activation during response inhibition, on amygdalar activation in response to fearful faces, on temporal activation during an auditory task, and on occipital activation during visual processing [13]. A third experiment in the same study showed that pretreatment with CBD prevented acute induction of THC-induced psychotic symptoms compared to pretreatment with placebo, and suggested that THC and CBD can have opposing effects on regionally-specific brain activation, which may underlie their different symptomatic and behavioural effects. Thus, there is converging *in vivo* evidence supporting a differential neurobiological effect for different cannabinoids in the brain.

The present critical review aims at integrating the above evidence with findings from animal models. We will specifically focus on rodent studies that have been used to improve our understanding of the pro-psychotic actions of THC and the anti-psychotic actions of CBD. Furthermore, we will carefully overview research that addresses interactions between these major cannabinoid constituents.

*Address correspondence to this author at the Discipline of Pharmacology, University of Sydney NSW 2006, Australia; Tel: +61 2 9351 6954; Fax: +61 2 9351 2658; E-mail: jonathon.arnold@sydney.edu.au

We will then discuss how the study of cannabinoid action in animal models of schizophrenia may provide a deeper understanding of the underlying mechanisms responsible for cannabis-induced psychosis and provide insight on how cannabinoid-based therapies might be utilised to treat this chronic, disabling mental disorder.

CANNABIS, CANNABINOIDS AND THE ENDOCANNABINOID SYSTEM

Cannabis contains more than 60 molecules that belong to the class known as cannabinoids. The science of cannabis and the cannabinoids has progressed dramatically in the last two decades. An important factor in this has been the explosion of knowledge of the endocannabinoid system that exists within our brain and body. Cannabinoid CB1 and CB2 receptors have been cloned and their distribution in the brain and periphery mapped out [16, 17]. Further endocannabinoid agonists were discovered with the most characterized being arachidonic acid derivatives anandamide and 2-arachidonoylglycerol (2-AG) [18, 19]. The enzymes responsible for the biosynthesis of anandamide require complete characterization but a N-acyl phosphatidylethanolamine phospholipase D (NAPE-PLD) has been implicated in the process [20]. The synthesis of 2-AG requires diacylglycerol lipase- α (DAGL- α) [21]. Metabolism of anandamide and 2-AG require fatty amide acid hydrolase (FAAH) and monoacylglycerol lipase (MAGL) respectively [20]. In neurons both anandamide and 2-AG behave as retrograde transmitters travelling backwards from the postsynaptic membrane to activate largely presynaptically located CB1 receptors to modulate the release of various neurotransmitters such as γ -aminobutyric acid (GABA), glutamate, dopamine and serotonin [22]. They also mediate synaptic plasticity such as depolarization-induced suppression of inhibition [23]. These aspects of the cannabinoid system are fully described elsewhere in this issue.

The main psychoactive constituent of cannabis, THC, is a partial agonist at both CB1 and CB2 receptors. THC through its interaction with the CB1 receptor found on neurons is largely responsible for the distinctive effects of cannabis on mood, cognition and behaviour. CB2 receptors are expressed on microglia (the brain's macrophages) and perhaps even neurons, which opens up a whole new layer of complexity in the understanding of cannabinoid neuropsychopharmacology [16]. Furthermore, some of the actions of THC have been attributed to interfering with endocannabinoid-mediated synaptic plasticity [24, 25]. Although THC is the most abundant cannabinoid in the plant, another component, CBD, has also been extensively characterized but its mechanisms are more opaque. CBD is an isomer of THC but the subtle distinction in their chemical structures confers marked differences in their pharmacological activities. Early studies showed CBD had little affinity for cannabinoid receptors, likely explaining its lack of psychoactivity [26]. However, recent evidence suggests it may be an antagonist/inverse agonist at CB1 and CB2 receptors [27]. CBD also activates a myriad of receptor proteins dependent on concentration including transient receptor potential vanilloid type 1 (TRPV1) and 5-HT1A receptors [26]. Furthermore CBD inhibits FAAH, enhancing the levels of anandamide at the synapse.

RODENT MODELS OF SCHIZOPHRENIA: METHODOLOGICAL CONSIDERATIONS

It is impossible to model the uniquely human condition of schizophrenia in its entire complexity using rodent models. At best these models approximate aspects of the disorder and provide a means to improve our understanding of neurobiological processes that are related to schizophrenia symptoms. In addition they provide a platform to screen new therapeutic treatments for the disorder. At the core of compelling rodent models of schizophrenia are measures of behavioural deficits in animals that are also observed in schizophrenia patients. For example schizophrenia patients display deficits in various cognitive functions, with the most robust being in

working memory [28]. Further, these patients have impaired sensorimotor gating as operationalised by prepulse inhibition (PPI) of startle or auditory evoked potentials [29-35]. All of these aforementioned phenomena have been reproduced in rodents. Negative symptoms like social withdrawal can also be measured in animals. Positive symptoms of the disorder such as hallucinations and delusions are - at least currently - impossible to measure in rodents. These symptoms are likely subserved by dopamine hyperactivity or glutamatergic hypoactivity in the brain of schizophrenia patients. Agents that increase dopamine levels or block N-methyl-D-aspartate (NMDA) receptors in the CNS trigger locomotor hyperactivity in rodents and so this behaviour is used to assess whether certain agents have antipsychotic potential by reversing such hyperactivity.

Schizophrenia arises due to genetic and environmental factors combining to negatively affect brain maturation at critical, early periods of development [36]. The translational power of rodent models will improve with the introduction of novel paradigms that combine genetic, environmental and neurodevelopmental factors to more faithfully approximate the underlying pathophysiology of the disorder. Current models are largely unidimensional, only recreating aspects of the disorder from single genetic or environmental factors that are related to schizophrenia [37]. Traditional rodent models of schizophrenia are based on dopamine hyperactivity and glutamate hypofunctioning theories of schizophrenia neurochemistry, replicating human evidence suggesting these are the core neurochemical alterations in the illness [38-40]. Agents that enhance dopamine (e.g. psychostimulants) or inhibit glutamate functioning in the CNS (NMDA receptor antagonists) trigger schizophrenia-like behaviours both in humans and animals. Sensitization of mesocorticolimbic dopaminergic circuits has also been suggested as important in the misattribution of salience to environmental stimuli, and may be involved in the positive symptoms of schizophrenia since its early phases [2, 38, 41, 42]. Therefore, rodent studies measuring behavioural and neurochemical sensitization to repeated psychoactive drug exposure have relevance to improving our understanding of the neurobiology of schizophrenia [43-45]. Early life stress has also been linked to increased risk of developing schizophrenia and animals exposed to neonatal stress (e.g. maternal deprivation or social isolation) display long-lasting schizophrenia-relevant deficits expressed in adulthood. With increasing knowledge of the genetic basis of schizophrenia newer genetic models have been developed where mutations in schizophrenia susceptibility genes [e.g. neuregulin 1 (*NRG1*)] promote schizophrenia-related behaviours. Many of these animal models have been used to examine the neural and genetic basis of the pro-psychotic and anti-psychotic potential of THC and CBD.

The current review will specifically address rodent studies that examine the effects of cannabinoids: on schizophrenia-related neurobehavioural phenotypes; in adolescence; and in dopamine, glutamate, neonatal stress and genetic models of schizophrenia (for a summary of the effects of THC and CBD in these models see Tables 1 and 2 respectively). We will also assess rodent evidence of the potential therapeutic efficacy of cannabinoid receptor antagonist drugs as novel antipsychotics, as well as evaluating whether cannabinoid receptor knockout mice display schizophrenia-relevant phenotypes. The final section of this review will be devoted to rodent studies addressing interactions between THC and CBD that might provide mechanistic insight into interactions between these agents that has been reported in the human literature (see Table 3).

EFFECTS OF CANNABINOIDS ON SCHIZOPHRENIA-RELATED NEUROBEHAVIOURAL PHENOTYPES IN RODENTS

Acute THC administration transiently exacerbates core psychotic symptoms and cognitive deficits in schizophrenia patients

Table 1. The Effects of THC and Cannabinoid (CB) Agonists in Rodent Models of Schizophrenia

Model	CB	Species	CB Effect	Ref.
Dopamine	THC (adult, repeated dosing)	Rat	Potentiated ↑ locomotor activity and stereotypy (amphetamine)	[119, 120]
	THC and CP 55,940 (adult, repeated dosing)	Rat Mouse	No effect on ↑ locomotor activity (cocaine)	[47, 82]
Glutamate	WIN 55,212-2 (adult, acute dosing)	Rat	Exacerbated ↓ short-term memory deficits and hippo- campal pyramidal cell firing (MK-801)	[133]
	THC (adolescent, repeated dosing)	Rat	Exacerbated ↓ recognition memory and GTPγS binding Blocked ↑ 2-AG levels in the prefrontal cortex (phenylclidine)	[134]
	WIN 55,212-2 (adult, repeated dosing)	Rat	Reversed ↓ PPI, recognition memory and social behav- iour (phenylclidine)	[135]
Neonatal stress	THC (adult, acute dosing)	Rat	Potentiated ↓ PPI	[68]
	CP 55,940 (adolescent, re- peated dosing)	Rat	Stress or CB alone ↓ CB1 receptor expression and ↑ astrocytes no. in the hippocampus – combined stress and CB nullified these effects	[154]
	CP 55,940 (adolescent, re- peated dosing)	Rat	No effect	[156]
	WIN 55,212 (neonatal, acute dosing)	Rat	Potentiated ↓ immune function Stress reversed and potentiated ↑ anxiety and serum corticosterone promoted by CB respectively (males only)	[157]
	THC (adult, acute dosing)	<i>Nrg1</i> HET mice	Enhanced behavioural effects of THC in <i>Nrg1</i> HET mice (e.g. <i>Nrg1</i> -specific ↑ PPI) <i>Nrg1</i> -specific ↑ brain activation pattern in stress cir- cuitry (e.g. LSV)	[57, 172, 175]
	CP 55,940 (adult, repeated dosing)	<i>Nrg1</i> HET mice	<i>Nrg1</i> modulated CB tolerance; ↑ FosB/ΔFosB expres- sion in LSV	[58]
	THC (adolescent, repeated dosing)	<i>Nrg1</i> HET mice	Exacerbated ↑ locomotor activity in <i>Nrg1</i> HET mice Promoted <i>Nrg1</i> -specific effects on CB1, 5-HT2A and NMDA receptor expression	[183]
	THC (adolescent, repeated dosing)	<i>COMT</i> KO mice	Induced <i>COMT</i> -specific ↑ locomotor activity, ↓ work- ing memory and ↓ anxiety	[186]
	WIN 55,212-2 (adolescent, repeated dosing)	<i>COMT</i> KO mice	Promoted <i>COMT</i> -specific ↓ PPI, ↑ social behaviour and ↓ anxiety	[187]
	THC (adult, acute dosing)	<i>Mdr1a/b</i> and <i>Bcrp1</i> KO mice	↑ hypothermia, and ↑ brain and blood [THC] in <i>Mdr1a/b</i> and <i>Bcrp1</i> KO mice.	[195, 197]

Table 2. The Effects of CBD In Rodent Models of Schizophrenia

Model	Species	CB Effect	Ref.
Dopamine	Mouse (adult, repeated dosing)	Blocked ↑ locomotor activity (amphetamine)	[60]
	Mouse Rat (adult, acute dosing)	Blocked ↑ locomotor activity (amphetamine)	[123, 124]
	Rat (adult, repeated dosing)	No effect on ↑ locomotor activity (amphetamine)	[125]
Glutamate	Mouse (adult, acute dosing)	Inhibited ↑ locomotor activity (ketamine)	[123]
	Mouse (adult, acute dosing)	Reversed ↓PPI (MK-801)	[136]
	Rat (adult, acute dosing)	No effect on ↓PPI Blunted social withdrawal	[137]
Genetic	<i>Nrg1</i> HET mice (adult, repeated dosing)	No effect on ↓PPI, ↑ locomotor activity and ↓ 5-HT _{2A} receptor expression in <i>Nrg1</i> HET mice. <i>Nrg1</i> -specific ↑ in social behaviour and GABA _A receptor expression.	[198]

Table 3. Rodent Studies Examining Interactions between THC and CBD

	CB Effect	Species	CBD: THC Ratio	Proposed Mechanism	Admin Schedule	Ref.
CBD ↑ effect of THC	Antinociception	Mouse	100:1 (30 mg/kg CBD 0.3 mg/kg THC)	CBD ↑ blood but not brain [THC]	CBD pretreatment (10 min) i.v.	[101]
	↓ Spatial memory Hypothermia	Mouse	50:1 (50 mg/kg CBD 1 mg/kg THC)	↑ CB ₁ receptor in hippocampus and hypothalamus	Co-administration (simultaneous) i.p.	[90]
	↓ Locomotor activity	Mouse	10:1 & 50:1	↑ CB ₁ receptor in hippocampus and hypothalamus	Co-administration (simultaneous) i.p.	[90]
	↓ Social behaviour ↓ Locomotor activity	Rat	2:1 (20 mg/kg CBD 10 mg/kg THC) NDD	Not examined	CBD pretreatment (20 min) i.p.	[61]
	Weight loss ↑ Anxiety ↓ Social behaviour	Rat (adolescent)	1:1 (1-10 mg/kg CBD & THC)	CBD ↑ brain & blood [THC] No effect on CB ₁ or 5-HT _{1A} receptors.	CBD pretreatment (20 min) i.p., repeated, escalating	[246]
CBD ↓ effect of THC	Antinociception	Mouse	3.3:1 (10 mg/kg CBD 3 mg/kg THC) NDD	Not examined	CBD pretreatment (10 min) i.v.	[101]
	↓ Spatial working memory	Rat	12.5:1 (50 mg/kg CBD 4 mg/kg THC)	Not examined	Co-administration (simultaneous) i.p.	[89]
	↓ Social behaviour	Rat	20:1 (20 mg/kg CBD 1 mg/kg THC)	Not examined	CBD pretreatment (20 min) i.p.	[61]
	Place aversion	Mouse	1:10 & 1:1 (1 & 10 mg/kg CBD 10 mg/kg THC) NDD	Not examined	Co-administration (simultaneous) i.p.	[241]

NDD = non dose-dependent

[7]. Furthermore, THC administered to healthy participants mimics some of the core symptoms of schizophrenia – increasing positive and negative schizophrenia-like symptoms and cognitive deficits including impairment in working memory [4]. Similarly in rodents THC administration promotes an array of effects that are thought relevant to aspects of schizophrenia symptomology. Locomotor hyperactivity, presumably due to increased dopamine release in the brain, has been reported for THC and its synthetic analogues but only when using relatively low doses of these drugs [46–49]. Such locomotor activating effects of THC are consistent with studies showing that acute exposure to cannabinoids promote dopamine release in the nucleus accumbens and medial prefrontal cortex ([50–53] for a good review of this work see [54]). At high doses THC promotes dose-dependent locomotor suppression in rodents and catalepsy due to dense distribution of CB1 receptors in motor-related regions of the brain such as the globus pallidus, substantia nigra and cerebellum [55, 56].

Acute administration of THC and THC-like cannabinoid receptor agonists (e.g. CP 55,940 and WIN 55,212-2) also promotes deficits in behaviours relevant to negative symptoms of schizophrenia such as social withdrawal. Exposure to cannabinoids reduces social interaction of rodents following acute or repeated exposure which can be dissociated from the locomotor suppressant effects of these drugs [57–63]. Endocannabinoid neurotransmission may be involved in the mediation of prosocial behaviour as, unlike the administration of cannabinoid receptor agonists, the inhibitor of FAAH URB597 increased social interaction in adolescent rats [64]. Indeed the balance between social approach and withdrawal appears differentially regulated by CB1 receptors localised on GABAergic versus glutamatergic neurons. That is, CB1 receptor deletion from cortical and striatal GABA interneurons enhanced social and novel object exploration, whereas when CB1 receptors were removed from cortical glutamatergic neurons social and object exploration was reduced [65].

Acute administration of cannabinoid receptor agonists to rodents promotes a number of impairments in attention and cognitive function that are relevant to schizophrenia (see the specific paper released in the present issue). Cannabinoid receptor agonists including THC promote PPI deficits in rodents that can be reversed by atypical antipsychotic drugs [58, 66, 67]. However, only one study was able to show an acute cannabinoid-induced PPI deficit without the confounding influence of these agents on acoustic startle [67]. Other studies have shown that cannabinoid receptor agonists may augment PPI or have no effect [60, 68, 69]. Synthetic cannabinoid receptor agonists also disturb auditory evoked potentials. CP 55,940 disrupted both auditory evoked potentials and theta oscillations in the CA3 region of the hippocampus and entorhinal cortex that were mediated by CB1 receptors [70]. Further, CB1 receptor-mediated disruption of auditory gating by WIN 55,212 has also been demonstrated in the CA3, dentate gyrus and medial prefrontal cortex [71].

Acute systemic THC administration decreased working memory performance that correlated with reduced acetylcholine release in the hippocampus that were reversed by antipsychotic drugs [72, 73]. CB1 receptors in the hippocampus are vital to the memory impairing effects of THC as intrahippocampal injection of the CB1 receptor antagonist SR 141716 reversed working memory deficits of systemically administered THC [74]. Injection of THC directly into the medial prefrontal cortex also disrupts spatial working memory, which was reversed by the dopaminergic D1 receptor antagonist SCH23390 and the atypical antipsychotic drug clozapine [75]. The amnesic effects of THC in the hippocampus involve CB1 receptors localised on GABA interneurons that activate the mTOR pathway leading to a maladaptive increase in protein translation [76]. In addition, THC impairs cognitive flexibility and auditory discrimination performance that is mediated by CB1 receptors in rats [77]. Interestingly spatial learning impairments and cognitive

inflexibility promoted by THC, unlike other effects of cannabinoids, are resistant to tolerance [78].

Behavioural sensitization is subserved by neurochemical sensitization in mesocorticolimbic circuits that might mediate the positive symptoms of schizophrenia [2, 41]. Behavioural sensitization is best characterized with repeated, intermittent dosing with psychostimulants where the initial enhancement in locomotion promoted by these drugs is progressively increased with repeated dosing. Early demonstrations of such behavioural sensitization to THC are questionable based on limitations in the experimental design implemented in these studies including a lack of tracing progressive changes in locomotor activity over days [79, 80]. The common design was to repeatedly administer vehicle or escalating THC doses and then challenge all animals with THC on a final test day 20 days after final pretreatment exposure [79, 80]. Arguing for “sensitization” the THC-challenged animals in the THC group had significantly higher locomotor activity or stereotyped behaviour scores than animals in the vehicle-pretreated group. However these studies lacked control groups critical to discerning sensitization (reverse tolerance) from tolerance. Without demonstrating that the THC-pretreatment group had significantly higher activity than a vehicle-vehicle or THC-vehicle group, the results reported might simply indicate lack of tolerance to the suppressant effects of THC in the vehicle-pretreated group compared to complete tolerance in the THC-pretreated animals. One more recent study implemented an identical test design to these early studies, included the critical experimental controls, and used both measures of locomotor activity and stereotyped behaviours. This study failed to demonstrate behavioural sensitization to THC [81]. Such findings are consistent with other work using designs typically utilised to demonstrate behavioural sensitization to psychostimulants where no such behavioural sensitization to cannabinoids could be demonstrated in animals of various age ranges [82–86].

While sensitization to repeated THC exposure is not expressed behaviourally it may be neurochemically due to various subtle neuroadaptive processes in the mesolimbic dopamine pathway. Repeated THC pretreatment reduced dopamine release in the shell and increased dopamine release in the core of the nucleus accumbens in response to THC challenge compared to animals pretreated with vehicle [87]. Repeated THC exposure also enhanced expression of Δ FosB in the nucleus accumbens and prefrontal cortex suggesting the drug triggers long-term neuroadaptive changes in mesocorticolimbic circuits [88]. In addition, repeated THC exposure promoted structural changes in neuronal morphology (increased dendritic branching and dendritic length) in medium spiny cells of the nucleus accumbens shell and in pyramidal cells of the medial prefrontal cortex [84]. Similar changes in Δ FosB and dendritic morphology have been observed following psychostimulant sensitization. Future studies are needed to better characterise the sensitizing neurobiological effects that repeated THC exposure might have on the mesocorticolimbic system.

Relative to THC, CBD appears largely behaviourally inert. CBD doesn't modulate social behaviour or learning and memory function in rodents following acute or repeated administration [60, 63, 89–92]. However, acute CBD exposure has anxiolytic effects in rodents that are mediated by 5-HT_{1A} receptors and TRPV1 receptors [93–99]. Repeated CBD dosing promoted non dose-related anxiolytic effects in mice [60]. Contrasting with this a recent paper reported that repeated CBD dosing in rats promoted an anxiogenic effect [100]. Of particular relevance to schizophrenia is the finding that acute and repeated CBD administration enhances PPI in mice [60]. Acute CBD promoted a dose-related PPI facilitation (1 – 50 mg/kg), however following repeated exposure it was only the 1 mg/kg dose that retained the ability to facilitate PPI. This suggests that lower doses might be preferable to avoid tolerance to CBD-induced PPI facilitation. CBD does not modulate locomotor activity in male C57BL/6JArc mice following either acute or repeated in-

Table 4. Outstanding Research Questions

Examination of the effects of cannabinoids in novel multidimensional animals models of schizophrenia which reflect genetic, environmental and neurodevelopmental aspects of the disorder.
Better characterization of the sensitizing neurobiological effects that repeated THC exposure has on the mesocorticolimbic system.
Elucidation of the mechanism of action of CBD's antipsychotic efficacy in various animal models of schizophrenia.
Isolation of further genes involved in moderating cannabinoid action on schizophrenia-related phenotypes.
Further characterization of the role of CB1 and CB2 receptors in schizophrenia-relevant phenotypes.
Reconciliation of why in some cases CBD potentiates the effects of THC, while in others it inhibits THC's action.

traperitoneal dosing across a wide dose-range [60]. Although biphasic effects of intravenous CBD in ICR mice have been reported with locomotor stimulation at 10 mg/kg and suppression at 30 mg/kg [101]. It has been argued that because CBD doesn't promote catalepsy it might be better tolerated in patients than current prescribed therapies, which do have this sedative effect in rodents. It has also been suggested that CBD might engender neuropharmacological effects akin to the atypical antipsychotic clozapine rather than the typical drug haloperidol, as CBD and clozapine administration increased c-Fos expression, a marker of brain activation, only in the ventral striatum whereas haloperidol promoted c-Fos expression in both the ventral and dorsal striatum [102].

ADOLESCENCE – A PERIOD OF VULNERABILITY TO CANNABINOID EXPOSURE

Adolescence is a significant period of neurodevelopment in the pathophysiology of schizophrenia coinciding with the onset of the disorder and first experimentation with cannabis (for an excellent review see [103]). Animal studies have shown that this period offers particular vulnerability to the actions of cannabinoids leading to long-term neurobehavioural deficits in adulthood. Pubertal exposure to the synthetic cannabinoid receptor agonist WIN 55,212-2 promoted long-lasting deficits in PPI, social interaction, and social and object recognition that were reversed by treatment with antipsychotic drugs such as haloperidol and quetiapine [104-107]. Adolescent rats are less sensitive to the aversive effects of cannabinoids than adults, however they are more sensitive to the detrimental effects of cannabinoid receptor agonists including THC on object recognition memory and working memory [108-111]. Taken together, this suggests adolescent cannabis use provides a dangerous scenario where users may be less averse to the immediate effects of the drug and more likely to use it repeatedly thus exposing themselves to long-term cognitive impairment.

The neural basis of altered vulnerability to the effects of cannabinoids in adolescence shows that the adolescent brain may be more or less sensitive to the effects of cannabinoids than the adult brain, dependent on the proteins measured or the brain region analysed. Exposure to the potent synthetic analogue of THC, HU-210, has less effects on the adolescent brain than the adult CNS in terms of dopamine D2, GABAA, 5-HT1A and CB1 receptor expression in various brain regions [112-114]. Although a single dose of HU-210 was sufficient to promote increases in dopamine D1 receptor expression in nigrostriatal and mesolimbic brain regions of adolescent rats, this was only attained in adult animals after repeated dosing [115]. During adolescence there exist dynamic changes in the levels of endocannabinoids and CB1 expression level in the prefrontal

cortex and nucleus accumbens, and repeated THC exposure enhanced anandamide levels in the nucleus accumbens selectively in adolescent animals [116].

Repeated THC exposure altered a greater number of proteins in the hippocampus of adolescent compared to adult rats (27 versus 10 respectively) [109]. Of note, proteins related to oxidative stress and protein folding were altered which could be related to greater degenerative changes and enhanced free radical damage in the hippocampus of adolescent animals. In addition, cytoskeletal and structural proteins were also affected indicating greater cytoarchitectural changes in the adolescent hippocampus. THC exposure selectively decreased levels of ubiquitin-conjugating enzyme E2 in adolescent rats. Interestingly, reduced levels of ubiquitin-conjugating enzymes have also been observed in the hippocampus of schizophrenia patients [117]. Repeated adolescent THC exposure also altered hippocampal dendritic morphology as well as reducing a number of synaptic markers including PSD-95 and expression levels of NMDA receptors [110]. Taken together hippocampal changes resulting from THC exposure in adolescence may contribute to lasting cognitive deficits that may have relevance to the pathophysiology of schizophrenia.

THE EFFECTS OF CANNABINOID IN DOPAMINE MODELS OF SCHIZOPHRENIA

Repeated THC treatment enhances the acute stimulatory effects of amphetamine and high doses of quinpirole (a D2 receptor agonist) on locomotor activity and stereotypy [118-120], suggesting that THC exposure sensitizes underlying mesocorticolimbic dopamine circuits. Prior studies have failed to demonstrate that THC or a 1:1 combination of THC and CBD modulates the hyperlocomotor effects of cocaine which is consistent with earlier studies utilising the cannabinoid receptor agonist CP 55,940 [47, 82]. Amphetamine promotes much greater levels of dopamine in the nucleus accumbens than cocaine [121, 122], therefore it is possible that only the action of potent psychostimulants like amphetamine is able to unmask subtle sensitizing changes in the mesocorticolimbic pathway triggered by repeated cannabinoid treatment.

The effect of CBD on neurobehavioural responses to psychostimulant drugs has been less comprehensively studied than that for THC and cannabinoid receptor agonists. Neuroadaptive changes in response to repeated CBD exposure might be necessary for demonstrating an antipsychotic action in dopamine models as repeated but not acute exposure to CBD (50 mg/kg) was effective in reversing amphetamine-induced hyperactivity in mice [60]. However, earlier studies have shown that acute CBD dose-dependently reduced amphetamine-induced hyperactivity in both mice and rats

[123, 124]. Another more recent study suggests that repeated CBD doesn't modulate locomotor hyperactivity in response to repeated exposure to amphetamine and behavioural sensitization [125]. This study utilized a study design that has been argued to model mania [126-128]. Two different treatment regimens were utilised in this study: 1) a "reversal" model where rats were pretreated with amphetamine twice daily for 7 days before CBD (15 – 60 mg/kg) was co-administered with amphetamine (2 mg/kg) on days 8-14 and 2) a "prevention" model rats where rats received CBD daily for 7 days before injection with amphetamine on days 8-14. In either case CBD did not influence the hyperlocomotor effects of amphetamine.

EFFECTS OF CANNABINOIDS IN GLUTAMATE MODELS OF SCHIZOPHRENIA

Phencyclidine, ketamine and MK-801, all glutamatergic NMDA receptor antagonists, have hallucinogenic effects in humans [129]. Administration of these agents to rodents promotes a number of different effects relevant to schizophrenia such as hypoactivation of the prefrontal cortex, deficits in PPI, cognitive dysfunction and social withdrawal [130-132]. Consistent with THC worsening cognitive deficits associated with schizophrenia, acute administration of WIN 55,212-2 exacerbated MK-801-induced deficits in short-term memory that correlated with an enhancement of the inhibitory effects of MK-801 on hippocampal pyramidal cell firing [133]. Another glutamate model utilized repeated, intermittent phencyclidine dosing over 4 weeks in adolescent rats before testing animals phencyclidine-free 3 days after the final dose. This schedule of phencyclidine treatment promoted a clozapine reversible impairment in recognition memory and alterations in the endocannabinoid system, notably in the prefrontal cortex, where phencyclidine reduced cannabinoid-stimulated GTP γ S binding and enhanced 2-AG levels [134]. Repeated co-administration of THC with phencyclidine in this model exacerbated impairments in recognition memory and the reduction in cannabinoid-stimulated GTP γ S binding but reversed the increase in 2-AG levels in the prefrontal cortex. Conflicting with this research, pre-exposure to the synthetic cannabinoid receptor agonist WIN 55,212-2 in adulthood reversed the usual deficits promoted by repeated phencyclidine such as PPI deficits, impaired recognition memory and social withdrawal [135]. Complicating interpretation of these data was that the "protective effects" of WIN 55,212-2 were on a background of the WIN 55,212-2 pre-treatment promoting deficits in PPI, recognition memory and social behaviour in control animals.

Unlike THC, CBD has an antipsychotic profile in glutamate models as it promoted a non dose-related inhibition of ketamine-induced hyperactivity (30 mg/kg was effective but not 15 or 60 mg/kg) in the absence of inhibiting motor activity on its own [123]. CBD also reversed acute MK-801-induced PPI deficits in mice [136] but not rats [137]. In addition, CBD blunted MK-801-induced social withdrawal in a non dose-dependent fashion as 3 and 10 mg/kg were effective, but not a higher 30 mg/kg dose [137]. The mechanism of action of CBD's antipsychotic action in glutamate models is unknown and its characterisation would be important to the development of this agent as an antipsychotic drug.

EFFECTS OF CANNABINOIDS ON NEONATAL STRESS MODELS OF SCHIZOPHRENIA

Early life stress is linked to an increased risk of psychosis, and childhood trauma may be involved in the pathogenesis of the disorder [138, 139]. Social isolation at weaning age has been used as an animal model of schizophrenia because it induces PPI deficits [140-142] and disrupts normal brain maturation [143, 144]. Acute THC exposure enhances PPI deficits promoted by post-weaning social isolation, an effect mediated by CB1 receptors [68]. Conflicting with this finding social isolation stress decreased CB1 receptor expression in the caudate putamen and amygdala [145]. It is possible that THC's exacerbation of social isolation-induced PPI deficits

is mediated by disruption of endocannabinoid tone in the brain. Indeed, social isolation stress increased FAAH expression in the caudate putamen and the nucleus accumbens [145], and expression of MAGL in the caudate putamen and DAGL α and DAGL β in the amygdala [146]. In addition, social isolation stress increased expression of NAPE-PLD and MAGL in the prefrontal cortex. Another stress-based and neurodevelopmental animal model of schizophrenia is the neonatal maternal stress deprivation model where rodents are separated from their mother at postnatal day 9 for 24 h. Adult animals that have undergone neonatal maternal deprivation display deficits in PPI, auditory sensory gating and startle habituation as well as reductions in levels of NR2A and NR2B NMDA receptor sub-units in the brain [147-151]. These animals display sexually dimorphic modulation of the endocannabinoid system – 13 day old male rats have increased levels of 2-AG and MAGL and decreased CB1 receptor expression in the hippocampus. Further both male and female animals display increased CB2 receptor and DAGL α expression in the hippocampus [152, 153].

Another promising approach combined two factors associated with increased risk of schizophrenia, neonatal stress and adolescent cannabinoid exposure, to observe whether there was any interaction following the sequential exposure to these factors [154]. Such an approach resonates with epidemiological evidence showing that childhood trauma and adolescent cannabis use increase the risk of schizophrenia by 3- and 2-fold respectively but when combined increase the risk by 21-fold [155]. Alone neonatal stress or repeated, adolescent exposure to the synthetic cannabinoid receptor agonist CP 55,940 reduced CB1 receptor expression and increased the number of GFAP-positive astrocytes in various regions of the hippocampus in male rats [154]. Somewhat surprisingly the combination of neonatal stress and adolescent cannabinoid exposure nullified these effects rather than potentiating them. Another study utilising the same paradigm observed no interactive effects on various schizophrenia-relevant behaviours in adulthood [156].

Neonatal maternal deprivation has protective effects against the anxiogenic effects of neonatal WIN 55,212-2 exposure in adolescent male but not female rats [157]. At odds with this finding is that neonatal maternal deprivation potentiated WIN 55,212-2-induced increases in serum corticosterone levels selectively in adolescent male rats [157]. This research shows that early neurodevelopmental stress and cannabinoid exposure may interact to modulate stress systems in the brain, that extends on prior research which has shown that the adult brain has a supra-additive response to stress and cannabinoid exposure [158-160]. Possibly as a consequence of altered HPA axis responsivity, neonatal stress and WIN 55,212-2 exposure additively reduced immune function as highlighted by reduced lymphocyte chemotaxis in the spleen and impaired lymphocyte proliferation in axillary nodes [157]. It would be of interest for future studies to continue with such an approach, possibly using other prenatal and neonatal stressors or immune challenges combined with adolescent exposure to the phytocannabinoid THC at doses more relevant to human consumption of cannabis.

GENETIC MOUSE MODELS OF VULNERABILITY TO CANNABIS-INDUCED SCHIZOPHRENIA

Genetic vulnerability to schizophrenia might explain why only a small proportion of the cannabis-using population develops schizophrenia. This is of great relevance to the clinical field of preventive psychiatry in human psychotic disorders. If the genes that confer genetic vulnerability to cannabis-induced psychosis were identified this would pave the way for preventative approaches where those at-high risk could be forewarned of the potential danger of experimentation with cannabis. In addition, one of the major impediments to the therapeutic development of cannabinoid medicines is the ability of these compounds to promote an increased risk of neuropsychiatric complications including psychosis. A pharmacogenetic approach would allow those at risk of psychiatric compli-

cations to be isolated leaving risk-free patients to benefit from cannabinoid-based therapy.

Early rodent studies attempting to model genetic vulnerability to the actions of cannabinoids compared the neurobehavioural responses of distinct inbred strains to cannabinoid exposure. Strain differences in the effects of cannabinoids were isolated on behaviour (e.g. reward and anxiety-related behaviours) [161, 162], brain activation patterns [163], and dopamine efflux in mesolimbic circuitry [52]. While these early studies lent credence to the idea that genetic disposition could moderate the actions of cannabinoids they did not provide any detail on the exact genes that were involved. The development of transgenic mice with targeted mutations has allowed identification of specific genes that moderate the CNS effects of cannabinoids, including those that modulate cannabinoid pharmacodynamics as well as the CNS disposition of these drugs.

We have utilized the heterozygous *Nrg1* mouse line (*Nrg1* HET mice) to develop an animal model of genetic vulnerability to cannabinoid-induced psychosis [164-167]. The neuregulin 1 gene (*NRG1*) - located on chromosome 8p12-p21 - was identified in 2002 as a schizophrenia-susceptibility gene [168]. The polypeptide influences key neurodevelopmental processes relevant to schizophrenia such as myelination, synaptogenesis, cell-cell signaling and neuronal migration, and regulates the expression/activation of NMDA receptors [169-171]. There are a number of ErbB receptors but ErbB4 is the only one that is bound and activated by NRG1 without forming a heterodimer. Alterations in the expression of *NRG1* mRNA have been found in the dorsolateral prefrontal cortex and hippocampus of patients with schizophrenia [169-171].

An initial study investigated the effects of THC on schizophrenia-relevant neurobehavioural parameters and demonstrated that adult male *Nrg1* mutant mice were more sensitive to the behavioural actions of THC compared to WT littermates [57]. The effects of THC on *Nrg1* HET mice were sex-specific as no increased behavioural sensitivity to acute THC challenge was detected in female *Nrg1* HET mice [172]. One striking finding was that unlike WT mice, male *Nrg1* HET mice showed improved PPI following THC exposure. Interestingly, nicotine and antipsychotic drugs increase PPI in rodents and reverse PPI deficits in patients with schizophrenia [173]. Following from our work recent human data support the idea that *NRG1* confers enhanced sensitivity to the actions of THC, however *NRG1* polymorphisms appear to worsen THC-induced information processing dysfunction rather than improve it. In healthy subjects THC-induced deficits in auditory mismatch negativity generation were only expressed in those with a specific *NRG1* variant [174]. These results suggest that variation in *NRG1* modulates CNS-effects of THC that can be related to dysfunctions observed in schizophrenia.

We also examined the neurobiological underpinnings of the increased behavioural susceptibility of *Nrg1* mutants to acute THC challenge by measuring THC-induced neuronal activation using c-Fos immunohistochemistry [175]. THC selectively increased c-Fos expression in the ventral part of the lateral septum (LSV) of *Nrg1* HET mice with no corresponding effect being observed in WT mice. Interestingly drugs which modulate sensorimotor gating, whether they are pro-psychotic drugs that impair PPI or antipsychotic drugs that facilitate PPI, all increase c-Fos expression in the lateral septum [176]. In addition, our study showed that THC promoted a greater increase in c-Fos expression in *Nrg1* mutants compared to WT mice in an array of stress-related regions of the brain such as the central nucleus of the amygdala and the paraventricular nucleus of the hypothalamus. A 3-way interaction between *Nrg1* hypomorphism, THC exposure and stress appeared necessary to this Fos expression profile as it was only observed in animals that underwent the stress of behavioural testing with no such effects being observed in animals administered THC in their homecage. Therefore, future studies examining *Nrg1*-stress-cannabinoid inter-

actions, particularly in a neurodevelopmental context, are warranted.

Chronic exposure to cannabinoids is more robustly associated with triggering psychosis than acute exposure so we examined how *Nrg1* HET mice responded to repeated cannabinoid administration [58]. Tolerance to CP 55,940-induced hypothermia and locomotor suppression developed more rapidly in adult, male *Nrg1* HET than WT mice. Conversely, only WT mice developed tolerance to the anxiogenic effect of CP 55,940 and *Nrg1* HET mice maintained consistent cannabinoid-induced anxiety throughout the repeated dosing period. Repeated cannabinoid exposure selectively increased FosB/ Δ FosB expression, a marker of long-term neuroadaptive changes, in the LSV of *Nrg1* HET but not WT mice. Opposite effects of acute CP 55,940 treatment were observed on sensorimotor gating, as PPI was facilitated in *Nrg1* hypomorphs and impaired in WT mice. Irrespective of genotype, tolerance developed to the acute effects of CP 55,940 on PPI with repeated exposure to the drug. These results highlight that *Nrg1* also modulates neuroadaptive responses to repeated cannabinoid exposure and further reinforces the notion that the LSV is an important brain region dysregulated in *Nrg1*-cannabinoid interactions.

The role of the LSV in *Nrg1*-cannabinoid interactions is not surprising given its role in the integration of cognitive and emotional information. The lateral septum receives cognitive input from the hippocampus as well as from the prefrontal cortex [177]. It also shares reciprocal projections with the hypothalamus and the amygdala that support its role in relaying affective information. When directly injected in the lateral septum, nicotine and serotonin receptor agonists induced anxiogenic effects in rats [178, 179]. The LSV predominantly expresses estrogen [177], which has been shown to modulate cannabinoid-induced presynaptic inhibition of glutamate and GABA release in the hypothalamus [180]. Further, estrogen receptor gene variants have been linked to increased risk of schizophrenia and estrogen receptors and ErbB4 interact in the transcriptional control of estrogen receptor target gene expression [181, 182]. Future studies addressing the involvement of estrogen in *Nrg1*-cannabinoid interactions in the LSV would be of interest, as well as studies determining the exact role of the LSV in mediating the behavioural effects of cannabinoids on *Nrg1* HET mice we have observed in our studies.

We have also examined whether *Nrg1* hypomorphism confers vulnerability to the actions of acute or repeated THC exposure in adolescence [183]. THC exposure in adolescence exacerbated the hyperlocomotor phenotype of *Nrg1* HET mice expressed after withdrawal of the drug. Furthermore, adolescent *Nrg1* HET mice were more resistant to THC-induced suppression of investigative social behaviours than WT mice. Repeated adolescent THC administration also promoted differential effects on CB1 receptor expression in the substantia nigra, with THC decreasing binding in WT mice while increasing it in *Nrg1* HET mice. The mRNA for ErbB4 receptors that bind Nrg1 is localized on dopaminergic neurons in the substantia nigra [184] therefore it is possible that CB1 and ErbB4 receptors work in concert in this brain region to regulate the hyperactive phenotype of *Nrg1* HET mice. *Nrg1* hypomorphism also altered the effects of adolescent THC exposure on neurotransmitter receptors implicated in the pathophysiology of schizophrenia, that is, 5-HT_{2A} and NMDA receptors. Notably in the insular cortex repeated adolescent THC exposure decreased 5-HT_{2A} receptor expression in WT mice but increased it in *Nrg1* mutants. Adolescent THC exposure also selectively increased NMDA receptor expression in the cingulate cortex and hippocampus or *Nrg1* HET but not WT mice.

The first gene implicated in conferring vulnerability to cannabis-induced psychosis was *COMT*, which encodes for an enzyme important in the degradation of dopamine in the CNS. Adolescent cannabis use significantly enhances the risk of experiencing positive symptoms and diagnosis with schizophreniform disorder in

those with a specific single nucleotide polymorphism in *COMT* [185]. Recent animal studies support the notion that *COMT* moderates the neurobehavioural actions of cannabinoids. *COMT* knockout mice treated with repeated THC in adolescence but not adulthood displayed sexually dimorphic (males only) locomotor hyperactivity, working memory deficits and increased anxiety-like behaviour unlike WT mice treated with THC [186]. In a follow-up study it was shown that *COMT* knockout mice were similarly more vulnerable to the neurobehavioural effects of repeated WIN 55,212-2 exposure in adolescence, and that the mechanism for this interaction didn't involve cannabinoid-modulation of the enhanced dopamine levels found in *COMT* knockout mice [187].

Apart from genes that vary the pharmacodynamic effects of cannabinoids on the brain, it is also possible that genes which regulate the CNS disposition of cannabinoids confer vulnerability to cannabis-induced psychosis. ABC transporters are drug efflux pumps expressed at various pharmacological barriers in the body and regulate the disposition of substrate drugs [188]. Cannabinoids have affinity for members of the ABC transporter superfamily such as multidrug resistance associated protein 1, breast cancer resistance protein and P-glycoprotein [189-192]. The most characterized ABC transporter is P-glycoprotein, which is expressed at the blood brain barrier and helps determine the CNS accumulation of substrate drugs including atypical antipsychotic drugs such as olanzapine and risperidone [193, 194]. Mice with a targeted deletion of the mouse homologue of the gene that encodes P-glycoprotein, *Mdr1a*, showed increased plasma concentrations of THC compared to WT mice [195]. Further, a *MDR1* variant increases the risk of developing cannabis dependence [196]. We have recently found that *Mdr1a/b* and *Bcrp1* knockout mice show impaired excretion of THC from the brain and display a prolonged THC-induced hypothermia than WT mice [197]. Therefore P-glycoprotein and breast cancer resistance protein regulate the CNS disposition of THC and examination of whether variation in *MDR1* or *BCRP* confers increased risk to cannabis-induced schizophrenia may provide a fruitful avenue of future research.

Thus far only one study has examined the antipsychotic efficacy of CBD in a genetic animal model of schizophrenia [198]. This study investigated whether: 1) *Nrg1* HET mice displayed an altered neurobehavioural response to CBD and; 2) if CBD reversed schizophrenia-related phenotypes expressed by these mice. Repeated CBD (1 – 100 mg/kg) was ineffective in reversing locomotor hyperactivity, PPI deficits and reduced 5-HT_{2A} receptor binding observed in *Nrg1* HET mice. However, repeated CBD (at 50 and 100 mg/kg) enhanced social interaction in *Nrg1* HET mice but not WT mice. Furthermore, acute CBD (100 mg/kg) selectively increased PPI in *Nrg1* HET mice, although tolerance to this effect was manifest upon repeated CBD exposure. Repeated CBD (50 mg/kg) also selectively increased GABA_A receptor binding in the granular retrosplenial cortex in *Nrg1* HET mice. These results suggest that *Nrg1* facilitates the neurobehavioural actions of CBD, but that CBD was unable to reverse schizophrenia-related behaviour exhibited by *Nrg1* HET mice. Future studies could examine whether adolescent CBD exposure is effective in reversing schizophrenia-related behaviour by hindering the manifestation of neurobehavioural disturbances displayed by *Nrg1* HET mice in adulthood.

THE EFFICACY OF CANNABINOID RECEPTOR ANTAGONISTS IN ANIMAL MODELS OF PSYCHOSIS

Cannabinoid CB1 receptor antagonists show some promise as novel antipsychotic drugs in animal models of schizophrenia. Pre-treatment with CB1 receptor antagonists SR141716 or AM251 reversed PPI deficits induced by acute apomorphine, phenylclidine and MK-801 in a similar fashion to the atypical antipsychotic drug clozapine [199]. This contrasted with an earlier finding where SR 141716 did not reverse PPI-deficits induced by apomorphine, am-

phetamine and MK-801 in rats [200]. These conflicting findings have been attributed to the different vehicle solutions, doses and PPI paradigms used between these studies. A novel cannabinoid CB1 receptor antagonist AVE1625 reversed MK-801 but not amphetamine induced abnormally persistent latent inhibition and improved working and recognition memory in mice [201]. This study suggested that AVE1625 might best be introduced as an adjunct therapy with conventional antipsychotic drugs because: 1) of its cognitive enhancing efficacy, 2) it did not alter the antipsychotic efficacy of haloperidol and olanzapine against phenylclidine and amphetamine-induced hyperactivity and 3) it ameliorated haloperidol and olanzapine-induced catalepsy and olanzapine-induced weight gain [201]. In a model where adolescent rats receive repeated, intermittent phenylclidine injections it was shown that co-administration with the CB1 receptor antagonist AM251 reversed phenylclidine-induced recognition memory deficits and enhancement of 2-AG levels in the prefrontal cortex [202].

Cannabinoid CB1 receptor antagonists inhibit behavioural sensitization promoted by psychostimulants [47, 203-206]. The literature contains conflicting findings on whether cannabinoid receptor antagonists hinder the induction versus expression of behavioural sensitization to psychostimulants that might be explained by the different cannabinoid CB1 receptor antagonists used in these studies and the differences in their selectivity [207]. AM 251 has been shown to impair induction [204-206] whereas SR 141716 has been shown to inhibit expression of behavioural sensitization to psychostimulants [47, 203]. Somewhat counterintuitive to these findings is that boosting endocannabinoid tone with URB 597, an inhibitor of FAAH, blocked the development of behavioural sensitization to amphetamine that was mediated by CB1 receptors [208]. The effectiveness of SR 141716 in reversing expression of behavioural sensitization to psychostimulants is also dependent on context, that is, SR 141716 only reversed established behavioural sensitization to cocaine when the drug was paired with the test chamber and not in mice receiving the drug in their homecage [47]. The VTA and nucleus accumbens have dissociable roles in mediating behavioural sensitization to psychostimulants, with the former being critical to induction and the latter to expression [209-211]. Interestingly, SR 141716 injected directly into the nucleus accumbens inhibited the expression but not the induction of behavioural sensitization to both cocaine [212] and methamphetamine [203]. Intraventricular SR 141716 injection was ineffective in reversing induction of behavioural sensitization to methamphetamine [203]. Methamphetamine-sensitized animals had reduced levels of CB1 receptors in the nucleus accumbens and these receptors were more sensitive to cannabinoid stimulation than animals treated with saline [203]. This further reinforces the notion that the endocannabinoid system is involved in contextual learning aspects of the expression of behavioural sensitization.

DO CANNABINOID RECEPTOR KNOCKOUT MICE DISPLAY SCHIZOPHRENIA-RELATED NEUROBEHAVIOURAL PHENOTYPES?

Human association studies have linked *CNR1*, the gene that encodes for the human cannabinoid receptor 1 gene, with an increased susceptibility to developing schizophrenia in various populations [213-216], although such positive associations have not always been replicated [217-219]. Lending weight to the notion that variation in *CNR1* is related to schizophrenia risk are observations that the brain of post-mortem schizophrenia patients contain increased CB1 receptor expression compared to healthy controls, particularly in areas of the brain that are distorted in schizophrenia patients such as the prefrontal cortex and cingulate cortex [220-223]. These studies suggest that increased CB1 expression or function may be involved in conferring susceptibility to schizophrenia.

CB1 receptor knockout mice do not display classic schizophrenia-like behaviour. Two separate CB1 receptor knockout mice have

been developed that were first described within months of each other [224, 225]. The CB1 receptor knockout mice developed by Ledent *et al.* (1999) bred on a CD1 background showed identical locomotor activity to WT animals, whereas the line developed by Zimmer *et al.* (1999) bred on a C57BL/6J background displayed locomotor hypoactivity. The Ledent line showed improved retention of novel object recognition memory function [226] and the Zimmer line exhibited disrupted reversal learning in the Morris water maze [227]. A follow up study confirmed that the deficit in reversal learning in these mice was due to an impairment in the extinction of spatial learning [228]. Consistent with the symptomology of schizophrenia, the Ledent line mice expressed reduced social behaviour dependent on being tested in an unfamiliar environment [229-231]. It would be of interest to observe whether mice overexpressing CB1 receptors provide a better animal model of schizophrenia given these findings with CB1 receptor knockout mice and that increased expression of the CB1 receptor is found in post-mortem schizophrenia brain [220-223].

It is possible that CB1 receptor loss might confer vulnerability to environmental factors that trigger psychosis in humans. CB1 receptor knockout mice display enhanced stereotypy promoted by phencyclidine administration consistent with an increased sensitivity to hyperdopaminergic function in these mice [230]. Consonant with this, SR141716 markedly increased stereotypies produced by simultaneous co-administration of both a D1 and D2 receptor agonists [232]. However, another study reported that CB1 receptor knockout mice were subsensitive to hyperlocomotion induced by cocaine and amphetamine [204]. In addition, CB1 receptor knockout hampered cocaine-induced c-Fos expression and phosphorylation of key proteins involved in the acute effects of cocaine in the striatum such as the GluR1 sub-unit of AMPA, DARPP-32 and extracellular-signal regulated kinase 2 (ERK2) [204]. CB1 receptors expressed in GABAergic medium spiny neurons of the striatum were critical to cocaine-induced ERK phosphorylation, which subserves the hyperlocomotor effects of cocaine. Research has also shown CB1 receptor deficiency blunted the actions of 5-HT1A and 5HT2A/C agonists due to impaired 5-HT1A and 5-HT2A/C receptor function [233]. It would be of interest to observe whether CB1 receptor knockout mice are more susceptible to the adverse neurobehavioural actions of other environmental challenges implicated in the aetiology of schizophrenia.

CB2 receptors were once believed to be expressed solely in the periphery, however, there is a growing body of evidence supporting their localisation in the CNS. CB2 receptors are expressed on microglia and there also exists controversial evidence suggesting their expression on neurons [16]. Two recent studies show that polymorphisms in *CNR2* increase the risk of developing schizophrenia and bipolar disorder [234, 235]. Two polymorphisms in CB2 receptors were more prevalent in two independent samples of schizophrenia patients [234]. The risk allele for one of these SNPs was associated with low levels of *CNR2* in human postmortem brain tissue. CHO cells transfected with a missense allele of one of the SNPs displayed impaired cannabinoid-induced reductions in cAMP levels. Mouse studies also showed that the CB2 receptor antagonist AM630 potentiated MK-801 and methamphetamine-induced PPI deficits and locomotor activity stimulation [234]. CB2 receptor knockout mice have recently been shown to display schizophrenia-related behaviours such as PPI deficits and impairments in short and long-term memory consolidation [236]. These animals, while exhibiting reduced baseline locomotor activity, were more sensitive to cocaine-induced hyperactivity compared to WT mice. Furthermore, PPI deficits in these mice were reversed by administration of the atypical antipsychotic risperidone. CB2 knockout mice also expressed greater D2 receptor and less 5-HT2A receptor mRNA expression in the prefrontal cortex than WT mice.

PRECLINICAL STUDIES ON INTERACTIONS BETWEEN THC AND CBD

THC levels in cannabis have been steadily increasing over the last three decades and the ratio of CBD:THC is decreasing [237-240]. A recent US potency study reported that between 1993 and 2008 THC concentrations in cannabis preparations have more than doubled while CBD content has remained the same [238]. For example in 1993 cannabis contained on average 0.3% CBD and 3.4% THC (1:11 CBD:THC ratio), whereas in 2008 it contained 0.4% CBD and 8.8% THC (1:22 ratio). One of the most striking changes was reported for hashish or cannabis resin which is normally assumed to have a 1:1 CBD:THC concentration. In 1993 hashish contained 3.8% CBD and 6.6% THC (1:1.7 ratio) but by 2008 THC concentrations dramatically increased and the CBD level approximately halved with 2.1% CBD and 23.1% THC (1:11 ratio). Interestingly, in the early 80's hashish contained approximately 6% CBD and 3% THC (2:1 ratio) [237].

Animal and human data suggest that THC is responsible for the psychosis-inducing effects of cannabis and that CBD may have antipsychotic potential. Thus, rising THC levels without any increase in CBD concentration is of major concern. To be sure that increasing CBD concentrations in cannabis will be of benefit, more research is needed to better understand the nature and mechanisms subserving THC-CBD interactions. Preclinical studies may provide the experimental control to unequivocally address this issue, however the existing literature contains much conflicting data. Some studies show that CBD reverses the pharmacological actions of THC [61, 89, 241-243], while others report potentiating effects [61, 90, 101, 244-246] or no effect at all [101, 247] (see Table 3).

One of most comprehensive studies on interactions between CBD and THC examined an array of dose combinations in the classic cannabinoid tetrad of tests in mice (locomotor activity, catalepsy, body temperature and nociception) [101]. Doses of CBD at a CBD:THC ratio between 30:1 and 1:1 had no effect on THC-induced locomotor suppression, catalepsy, hypothermia or antinociception. Only when the CBD:THC ratio was raised to 100:1 (ie 30 mg/kg CBD) was a potentiating action of CBD on antinociception promoted by a sub-threshold dose of THC realised (0.3 mg/kg). Interestingly the same 30 mg/kg CBD dose did not affect the antinociceptive actions of marijuana smoke suggesting other constituents in the plant might also modulate CBD-THC interactions. CBD (at 10 mg/kg but not 30 mg/kg) inhibited the analgesic actions of a maximal effective dose of THC (3.3:1 ratio). Taken together, pharmacological interactions between CBD and THC may not only rely on the ratio of THC:CBD used but also the actual doses of THC administered. It appears for any modulation to occur high doses of CBD are required and that low relative doses of THC favour potentiation and high doses antagonism. The fact that marijuana smoke also nullified THC-CBD interactions further highlights that a multitude of cannabinoid and non-cannabinoid constituents of the plant need to be taken into account when trying to simulate the complex pharmacology of cannabis use by humans.

Research has also attempted to address whether CBD modulates more complex psychological phenomena of relevance to schizophrenia and other mental disorders. Human users of cannabis low in CBD are more vulnerable to the memory impairing effects of THC [248]. Using THC-rich (high THC and low CBD and other phytocannabinoids) and CBD-rich extracts (high CBD and low THC etc), it was shown that CBD reversed the impairing effects of THC on spatial working memory as measured using a delayed-matching-to-place version of the open-field water maze in rats (only though if the ratio of CBD:THC was > 10:1; in this case 50 mg/kg CBD to 4 mg/kg THC) [89]. Conflicting with this finding using a more straightforward experimental design it was shown that 50 mg/kg CBD potentiated the impairing effects of 1 mg/kg THC on spatial memory as measured in an 8-arm radial arm maze in mice when the

drugs were co-administered simultaneously [90]. CBD also enhanced the hypothermic, cataleptic and motor suppressant actions of THC.

A number of recent studies have assessed whether CBD modulates social withdrawal and anxiety-related behaviour induced by THC. The social withdrawal induced by low dose THC (1 mg/kg) was reversed in animals pretreated with CBD (20 mg/kg administered 20 min before THC) [61]. However when a higher dose of THC (10 mg/kg) was administered significant social withdrawal and locomotor suppression was observed that was absent in animals treated with 10 mg/kg THC alone [61]. These findings conflict with the view that CBD is more likely to potentiate low THC doses and antagonise high ones, suggesting that CBD modulation of THC effects may vary from measure to measure. A more recent study assessed interactions between repeated co-administration of escalating doses of THC and CBD (CBD was administered 20 min before THC) on social behaviour and anxiety [246]. CBD enhanced the sedative and anxiogenic actions of THC including social withdrawal. It was also found that CBD exacerbated loss of body weight that is normally observed following repeated THC exposure [249].

Animal studies have investigated whether CBD modulates the subjective qualities of THC intoxication using place conditioning and drug discrimination paradigms. A particularly interesting observation that requires further investigation is that, while THC had no modulatory effect on place conditioning in adolescent rats, CBD and THC combined tended to promote a conditioned place preference [i.e. THC was more rewarding in the presence of CBD (1:1 ratio) [246]]. Using mice it was shown that CBD might hinder the aversive effects of THC on rodents as CBD (1 or 10 mg/kg but not 30 mg/kg) reversed conditioned place aversion induced by THC (10 mg/kg) (i.e. 1:10 and 1:1 ratios respectively) [241]. These findings are particularly important as they demonstrate interactions between CBD and THC at CBD:THC ratios relevant to those found in cannabis cultivated for recreational use today. The latter study used the drug discrimination procedure and also demonstrated that CBD doesn't modulate the subjective qualities of THC intoxication [241]. A prior study using much higher CBD:THC dosing ratios (30:1 and 100:1) showed CBD prolonged the discriminative stimulus properties of THC [245].

Interactions between CBD and THC are not adequately understood and may involve a multitude of both pharmacodynamic and pharmacokinetic mechanisms that require further investigation. CBD's ability to *hinder* THC effects might be more straightforwardly explained by CBD behaving as a potent antagonist at both CB1 and CB2 receptors [27]. Although CBD amelioration of THC effects often lacks dose dependence [61, 241] which might be due to non-selective effects of CBD being manifest at higher doses. CBD has a complex pharmacological profile and interacts with numerous non-cannabinoid receptor protein targets across a range of different concentrations (see [26]). For example, CBD activates 5-HT_{1A} receptors at relatively high concentrations [250]. Future studies are required to delineate the exact pharmacodynamic mechanisms responsible for CBD inhibition of THC action.

Evidence supporting both pharmacodynamic and pharmacokinetic explanations of CBD-induced *augmentation* of THC effects has also been provided. First, combined THC and CBD exposure increased CB1 receptor expression in the hippocampus and the hypothalamus that correlated with greater memory-impairing and hypothermic effects of THC respectively [90]. Contrary to this finding CBD co-administration did not modify repeated THC-induced reductions in CB1 receptor expression in a more recent study [246]. The latter study did, however, provide a pharmacokinetic explanation consistent with earlier studies [251-253] which have demonstrated that CBD treatment increases brain and blood levels of THC. These effects have been attributed to CBD inactivation of cytochrome P450 enzymes involved in the metabolism of THC [254-256]. The drug efflux pumps P-glycoprotein and breast

cancer resistance protein, which are expressed at the blood brain barrier, may also be involved as CBD inhibits these transporters [190, 192] and THC is a substrate for these transporters [195-197]. Thus, it is also possible that CBD blocks the excretion of THC from the brain.

CONCLUSION

Rodent studies assessing the actions of cannabinoids in animal models of schizophrenia have generated a wealth of information and a series of outstanding questions that could be addressed in future studies (see Table 4). Modeling the neurobiological pathophysiology responsible for cannabis-induced schizophrenia using rodents is limited by the validity of the models used to address this issue. The vast majority of studies have assessed the effects of cannabinoids in models that manipulate a single factor implicated in the aetiology of the disorder such as exposure to a particular environmental factor (e.g. psychostimulant exposure). A few promising studies have utilized environmental challenges at critical neurodevelopmental periods of development (e.g. neonatal and adolescent stages). The vast majority of these studies highlight that THC is a propsychotic agent that exacerbates schizophrenia-related behaviours and neurobiology in these models. These studies collectively show that CB1 receptors mediate the schizophrenia-related actions of THC, however, new evidence showing CB2 receptor knockout mice display schizophrenia-related behaviours implies that a role of CB2 receptors should also be considered in future research. Research highlights that adolescence, a time where individuals begin to experiment with cannabis and manifest schizophrenia symptoms, is a particular period of vulnerability to the actions of THC and its synthetic analogues. Cannabis triggers schizophrenia in only those with a genetic vulnerability to the disorder and in the last 5 years animal studies have begun to isolate genes that might be involved. Transgenic mice studies suggest that variation in the schizophrenia susceptibility gene *neuregulin 1* and the ABC transporter gene that encodes P-glycoprotein may contribute to cannabis-induced psychosis according to pharmacodynamic and pharmacokinetic mechanisms respectively. Further these studies reinforce human data demonstrating a role of *COMT* in this phenomenon. Future rodent studies could go one step further and examine the neurobehavioural consequences of combining genetic vulnerability (transgenic animals with mutations in schizophrenia susceptibility genes), with environmental challenges early in life (e.g. prenatal stress) and then adolescent cannabinoid exposure. Such an approach may more faithfully resemble the pathophysiological changes observed in schizophrenia patients and provide a better platform to test novel treatments.

While much research has been conducted assessing the propsychotic actions of THC and cannabinoid receptor agonists, few studies have assessed the antipsychotic potential of CBD and other novel cannabinoid-based therapies in animal models of schizophrenia. CBD does ameliorate schizophrenia-related behaviours induced by psychostimulant and NMDA receptor antagonist exposure, however the mechanisms responsible for CBD's antipsychotic profile have not been adequately explained. A repeated theme in many of these studies is that CBD's antipsychotic efficacy is not dose-dependent and in many cases follows an inverted U function. This likely reflects the complex pharmacology of CBD and its proclivity to affect many different proteins beyond simple CB1 receptor antagonism. Recent human data suggest that CBD hinders the propsychotic actions of THC and that the ratio of CBD:THC is decreasing in cannabis smoked by users. Rodent studies conducted over close to 40 years have attempted to model interactions between CBD and THC and have yielded more questions than providing unequivocal answers. Some of the literature reports that CBD reduces the pharmacological effects of THC, while other studies demonstrate the opposite, that CBD potentiates the actions of THC by increasing CB1 receptor expression levels in the brain and also augmenting THC concentrations in the brain and the blood. Future research is

needed to reconcile these discordant findings including studies that also assess modulating actions of other phytocannabinoids.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflicts of interest.

ACKNOWLEDGEMENTS

This work was supported by a University of Sydney Bridging Grant to JCA. TK is supported by a Career Development Award (568752) of the National Health and Medical Research Council of Australia (NHMRC) and a NHMRC project grant (1003886). TK is also supported by the Schizophrenia Research Institute utilizing infrastructure funding from NSW Health and by the Motor Neuron Disease Research Institute of Australia. TK and JCA are also supported by the Brain & Behavior Research Foundation (formerly National Alliance for Research on Schizophrenia and Depression: Young Investigator Awards). AAB is supported by a University of Sydney Postdoctoral Fellowship.

REFERENCES

- [1] Perala J, Suvisaari J, Saarni SI, *et al.* Lifetime prevalence of psychotic and bipolar I disorders in a general population. *Arch Gen Psychiatry* 2007; 64(1): 19-28.
- [2] van Os J, Kapur S. Schizophrenia. *Lancet* 2009; 374(9690): 635-45.
- [3] Tamminga CA. The neurobiology of cognition in schizophrenia. *J Clin Psychiatry* 2006; 67 Suppl 9: 9-13; discussion 36-42.
- [4] D'Souza DC, Perry E, MacDougall L, *et al.* The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. *Neuropsychopharmacology* 2004; 29(8): 1558-72.
- [5] Ramaekers JG, Kauert G, Theunissen EL, Toennes SW, Moeller MR. Neurocognitive performance during acute THC intoxication in heavy and occasional cannabis users. *J Psychopharmacol* 2009; 23(3): 266-77.
- [6] D'Souza DC. Cannabinoids and psychosis. *Int Rev Neurobiol* 2007; 78: 289-326.
- [7] D'Souza DC, Abi-Saab WM, Madonick S, *et al.* Delta-9-tetrahydrocannabinol effects in schizophrenia: implications for cognition, psychosis, and addiction. *Biol Psychiatry* 2005; 57(6): 594-608.
- [8] Bergamaschi MM, Queiroz RH, Chagas MH, *et al.* Cannabidiol reduces the anxiety induced by simulated public speaking in treatment-naïve social phobia patients. *Neuropsychopharmacology* 2011; 36(6): 1219-26.
- [9] Crippa JA, Derenusson GN, Ferrari TB, *et al.* Neural basis of anxiolytic effects of cannabidiol (CBD) in generalized social anxiety disorder: a preliminary report. *J Psychopharmacol* 2011; 25(1): 121-30.
- [10] Crippa JA, Zuardi AW, Martin-Santos R, *et al.* Cannabis and anxiety: a critical review of the evidence. *Hum Psychopharmacol* 2009; 24(7): 515-23.
- [11] Morgan CJ and Curran HV. Effects of cannabidiol on schizophrenia-like symptoms in people who use cannabis. *Br J Psychiatry* 2008; 192(4): 306-7.
- [12] Zuardi AW, Crippa JA, Hallak JE, Moreira FA, Guimaraes FS. Cannabidiol, a Cannabis sativa constituent, as an antipsychotic drug. *Braz J Med Biol Res* 2006; 39(4): 421-9.
- [13] Bhattacharyya S, Morrison PD, Fusar-Poli P, *et al.* Opposite effects of delta-9-tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology* 2010; 35(3): 764-74.
- [14] Fusar-Poli P, Crippa JA, Bhattacharyya S, *et al.* Distinct effects of {delta}9-tetrahydrocannabinol and cannabidiol on neural activation during emotional processing. *Arch Gen Psychiatry* 2009; 66(1): 95-105.
- [15] Fusar-Poli P, Allen P, Bhattacharyya S, *et al.* Modulation of effective connectivity during emotional processing by Delta 9-tetrahydrocannabinol and cannabidiol. *Int J Neuropsychopharmacol* 2010; 13(4): 421-32.
- [16] Atwood BK, Mackie K. CB2: a cannabinoid receptor with an identity crisis. *Br J Pharmacol* 2010; 160(3): 467-79.
- [17] Mackie K. Cannabinoid receptors: where they are and what they do. *J Neuroendocrinol* 2008; 20 Suppl 1: 10-4.
- [18] Devane WA, Hanus L, Breuer A, *et al.* Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* 1992; 258(5090): 1946-9.
- [19] Stella N, Schweitzer P, Piomelli D. A second endogenous cannabinoid that modulates long-term potentiation. *Nature* 1997; 388(6644): 773-8.
- [20] Howlett AC, Reggio PH, Childers SR, *et al.* Endocannabinoid tone versus constitutive activity of cannabinoid receptors. *Br J Pharmacol* 2011; 163(7): 1329-43.
- [21] Piomelli D. The molecular logic of endocannabinoid signalling. *Nat Rev Neurosci* 2003; 4(11): 873-84.
- [22] Freund TF, Katona I, Piomelli D. Role of endogenous cannabinoids in synaptic signaling. *Physiol Rev* 2003; 83(3): 1017-66.
- [23] Davies SN, Pertwee RG, Riedel G. Functions of cannabinoid receptors in the hippocampus. *Neuropharmacology* 2002; 42(8): 993-1007.
- [24] Mato S, Chevaleyre V, Robbe D, *et al.* A single in-vivo exposure to delta 9THC blocks endocannabinoid-mediated synaptic plasticity. *Nat Neurosci* 2004; 7(6): 585-6.
- [25] Mato S, Robbe D, Puente N, Grandes P, Manzoni OJ. Presynaptic homeostatic plasticity rescues long-term depression after chronic Delta 9-tetrahydrocannabinol exposure. *J Neurosci* 2005; 25(50): 11619-27.
- [26] Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol* 2008; 153(2): 199-215.
- [27] Thomas A, Baillie GL, Phillips AM, *et al.* Cannabidiol displays unexpectedly high potency as an antagonist of CB1 and CB2 receptor agonists *in vitro*. *Br J Pharmacol* 2007; 150(5): 613-23.
- [28] Kalkstein S, Hurford I, Gur RC. Neurocognition in schizophrenia. *Curr Top Behav Neurosci* 2010; 4: 373-90.
- [29] Braff DL, Geyer MA. Sensorimotor gating and schizophrenia. Human and animal model studies. *Arch Gen Psychiatry* 1990; 47(2): 181-8.
- [30] Braff DL, Geyer MA, Swerdlow NR. Human studies of prepulse inhibition of startle: normal subjects, patient groups, and pharmacological studies. *Psychopharmacology (Berl)* 2001; 156(2-3): 234-58.
- [31] Geyer MA. Are cross-species measures of sensorimotor gating useful for the discovery of procognitive cotreatments for schizophrenia? *Dialogues Clin Neurosci* 2006; 8(1): 9-16.
- [32] Hajos M. Targeting information-processing deficit in schizophrenia: a novel approach to psychotherapeutic drug discovery. *Trends Pharmacol Sci* 2006; 27(7): 391-8.
- [33] Powell SB, Zhou X, Geyer MA. Prepulse inhibition and genetic mouse models of schizophrenia. *Behav Brain Res* 2009; 204(2): 282-94.
- [34] Swerdlow NR, Braff DL, Geyer MA. Animal models of deficient sensorimotor gating: what we know, what we think we know, and what we hope to know soon. *Behav Pharmacol* 2000; 11(3-4): 185-204.
- [35] Turetsky BI, Calkins ME, Light GA, *et al.* Neurophysiological endophenotypes of schizophrenia: the viability of selected candidate measures. *Schizophr Bull* 2007; 33(1): 69-94.
- [36] Rapoport JL, Addington AM, Frangou S, Psych MR. The neurodevelopmental model of schizophrenia: update 2005. *Mol Psychiatry* 2005; 10(5): 434-49.
- [37] Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nat Neurosci* 2010; 13(10): 1161-9.
- [38] Fusar-Poli P, Howes OD, Allen P, *et al.* Abnormal prefrontal activation directly related to pre-synaptic striatal dopamine dysfunction in people at clinical high risk for psychosis. *Mol Psychiatry* 2011; 16(1): 67-75.
- [39] Fusar-Poli P, McGuire P, Borgwardt S. Mapping prodromal psychosis: A critical review of neuroimaging studies. *Eur Psychiatry* 2012; 27(3): 181-91.
- [40] Fusar-Poli P, Stone JM, Broome MR, *et al.* Thalamic glutamate levels as a predictor of cortical response during executive functioning in subjects at high risk for psychosis. *Arch Gen Psychiatry* 2011; 68(9): 881-90.
- [41] Featherstone RE, Kapur S, Fletcher PJ. The amphetamine-induced sensitized state as a model of schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2007; 31(8): 1556-71.

- [42] Fusar-Poli P, Howes OD, Allen P, *et al.* Abnormal frontostriatal interactions in people with prodromal signs of psychosis: a multimodal imaging study. *Arch Gen Psychiatry* 2010; 67(7): 683-91.
- [43] Tenn CC, Fletcher PJ, Kapur S. Amphetamine-sensitized animals show a sensorimotor gating and neurochemical abnormality similar to that of schizophrenia. *Schizophr Res* 2003; 64(2-3): 103-14.
- [44] Tenn CC, Fletcher PJ, Kapur S. A putative animal model of the "prodromal" state of schizophrenia. *Biol Psychiatry* 2005; 57(6): 586-93.
- [45] Tenn CC, Kapur S, Fletcher PJ. Sensitization to amphetamine, but not phencyclidine, disrupts prepulse inhibition and latent inhibition. *Psychopharmacology (Berl)* 2005; 180(2): 366-76.
- [46] Davis WM, Moreton JE, King WT, Pace HB. Marihuana on locomotor activity: biphasic effect and tolerance development. *Res Commun Chem Pathol Pharmacol* 1972; 3(1): 29-35.
- [47] Gerdeman GL, Schechter JB, French ED. Context-specific reversal of cocaine sensitization by the CB1 cannabinoid receptor antagonist rimonabant. *Neuropsychopharmacology* 2008; 33(11): 2747-59.
- [48] McGregor IS, Issakidis CN, Prior G. Aversive effects of the synthetic cannabinoid CP 55,940 in rats. *Pharmacol Biochem Behav* 1996; 53(3): 657-64.
- [49] Sanudo-Pena MC, Romero J, Seale GE, Fernandez-Ruiz JJ, Walker JM. Activational role of cannabinoids on movement. *Eur J Pharmacol* 2000; 391(3): 269-74.
- [50] Chen J, Paredes W, Lowinson JH, Gardner EL. Delta 9-tetrahydrocannabinol enhances presynaptic dopamine efflux in medial prefrontal cortex. *Eur J Pharmacol* 1990; 190(1-2): 259-62.
- [51] Chen JP, Paredes W, Li J, *et al.* Delta 9-tetrahydrocannabinol produces naloxone-blockable enhancement of presynaptic basal dopamine efflux in nucleus accumbens of conscious, freely-moving rats as measured by intracerebral microdialysis. *Psychopharmacology (Berl)* 1990; 102(2): 156-62.
- [52] Chen JP, Paredes W, Lowinson JH, Gardner EL. Strain-specific facilitation of dopamine efflux by delta 9-tetrahydrocannabinol in the nucleus accumbens of rat: an *in vivo* microdialysis study. *Neurosci Lett* 1991; 129(1): 136-80.
- [53] Tanda G, Pontieri FE, Di Chiara G. Cannabinoid and heroin activation of mesolimbic dopamine transmission by a common mu1 opioid receptor mechanism. *Science* 1997; 276(5321): 2048-50.
- [54] Kuepper R, Morrison PD, van Os J, *et al.* Does dopamine mediate the psychosis-inducing effects of cannabis? A review and integration of findings across disciplines. *Schizophr Res* 2010; 121(1-3): 107-17.
- [55] Herkenham M, Lynn AB, Johnson MR, *et al.* Characterization and localization of cannabinoid receptors in rat brain: a quantitative *in vitro* autoradiographic study. *J Neurosci* 1991; 11(2): 563-83.
- [56] Howlett AC, Bidaut-Russell M, Devane WA, *et al.* The cannabinoid receptor: biochemical, anatomical and behavioral characterization. *Trends Neurosci* 1990; 13(10): 420-3.
- [57] Boucher AA, Arnold JC, Duffy L, *et al.* Heterozygous neuregulin 1 mice are more sensitive to the behavioural effects of Delta(9)-tetrahydrocannabinol. *Psychopharmacology (Berl)* 2007; 192(3): 325-36.
- [58] Boucher AA, Hunt GE, Micheau J, *et al.* The schizophrenia susceptibility gene neuregulin 1 modulates tolerance to the effects of cannabinoids. *Int J Neuropsychopharmacol* 2011; 14(5): 631-43.
- [59] Genn RF, Tucci S, Marco EM, Viveros MP, File SE. Unconditioned and conditioned anxiogenic effects of the cannabinoid receptor agonist CP 55,940 in the social interaction test. *Pharmacol Biochem Behav* 2004; 77(3): 567-73.
- [60] Long LE, Chesworth R, Huang XF, *et al.* A behavioural comparison of acute and chronic Delta9-tetrahydrocannabinol and cannabidiol in C57BL/6JArc mice. *Int J Neuropsychopharmacol* 2010; 13(7): 861-76.
- [61] Malone DT, Jongejan D, Taylor DA. Cannabidiol reverses the reduction in social interaction produced by low dose Delta(9)-tetrahydrocannabinol in rats. *Pharmacol Biochem Behav* 2009; 93(2): 91-6.
- [62] O'Shea M, McGregor IS, Mallet PE. Repeated cannabinoid exposure during perinatal, adolescent or early adult ages produces similar longlasting deficits in object recognition and reduced social interaction in rats. *J Psychopharmacol* 2006; 20(5): 611-21.
- [63] van Ree JM, Niesink RJ, Nir I. delta 1-Tetrahydrocannabinol but not cannabidiol reduces contact and aggressive behavior of rats tested in dyadic encounters. *Psychopharmacology (Berl)* 1984; 84(4): 561-5.
- [64] Trezza V, Vanderschuren LJ. Bidirectional cannabinoid modulation of social behavior in adolescent rats. *Psychopharmacology (Berl)* 2008; 197(2): 217-27.
- [65] Haring M, Kaiser N, Monory K, Lutz B. Circuit specific functions of cannabinoid CB1 receptor in the balance of investigatory drive and exploration. *PLoS One* 2011; 6(11): e26617.
- [66] Nagai H, Egashira N, Sano K, *et al.* Antipsychotics improve Delta9-tetrahydrocannabinol-induced impairment of the prepulse inhibition of the startle reflex in mice. *Pharmacol Biochem Behav* 2006; 84(2): 330-6.
- [67] Schneider M and Koch M. The cannabinoid agonist WIN 55,212-2 reduces sensorimotor gating and recognition memory in rats. *Behav Pharmacol* 2002; 13(1): 29-37.
- [68] Malone DT, Taylor DA. The effect of Delta9-tetrahydrocannabinol on sensorimotor gating in socially isolated rats. *Behav Brain Res* 2006; 166(1): 101-9.
- [69] Stanley-Cary CC, Harris C, Martin-Iverson MT. Differing effects of the cannabinoid agonist, CP 55,940, in an alcohol or Tween 80 solvent, on prepulse inhibition of the acoustic startle reflex in the rat. *Behav Pharmacol* 2002; 13(1): 15-28.
- [70] Hajos M, Hoffmann WE, Kocsis B. Activation of cannabinoid-1 receptors disrupts sensory gating and neuronal oscillation: relevance to schizophrenia. *Biol Psychiatry* 2008; 63(11): 1075-83.
- [71] Dissanayake DW, Zachariou M, Marsden CA, Mason R. Auditory gating in rat hippocampus and medial prefrontal cortex: effect of the cannabinoid agonist WIN55,212-2. *Neuropharmacology* 2008; 55(8): 1397-404.
- [72] Egashira N, Ishigami N, Mishima K, *et al.* Delta9-Tetrahydrocannabinol-induced cognitive deficits are reversed by olanzapine but not haloperidol in rats. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32(2): 499-506.
- [73] Nava F, Carta G, Battasi AM, Gessa GL. D(2) dopamine receptors enable delta(9)-tetrahydrocannabinol induced memory impairment and reduction of hippocampal extracellular acetylcholine concentration. *Br J Pharmacol* 2000; 130(6): 1201-10.
- [74] Wise LE, Thorpe AJ, Lichtman AH. Hippocampal CB(1) receptors mediate the memory impairing effects of Delta(9)-tetrahydrocannabinol. *Neuropsychopharmacology* 2009; 34(9): 2072-80.
- [75] Rodrigues LC, Conti CL, Nakamura-Palacios EM. Clozapine and SCH 23390 prevent the spatial working memory disruption induced by Delta9-THC administration into the medial prefrontal cortex. *Brain Res* 2011; 1382: 230-7.
- [76] Puighermanal E, Marsicano G, Busquets-Garcia A, *et al.* Cannabinoid modulation of hippocampal long-term memory is mediated by mTOR signaling. *Nat Neurosci* 2009; 12(9): 1152-8.
- [77] Sokolic L, Long LE, Hunt GE, Arnold JC, McGregor IS. Disruptive effects of the prototypical cannabinoid Delta-tetrahydrocannabinol and the fatty acid amide inhibitor URB-597 on go/no-go auditory discrimination performance and olfactory reversal learning in rats. *Behav Pharmacol* 2011; 22(3): 191-202.
- [78] Boucher AA, Vivier L, Metna-Laurent M, *et al.* Chronic treatment with Delta(9)-tetrahydrocannabinol impairs spatial memory and reduces zif268 expression in the mouse forebrain. *Behav Pharmacol* 2009; 20(1): 45-55.
- [79] Cadoni C, Pisanu A, Solinas M, Acquas E, Di Chiara G. Behavioural sensitization after repeated exposure to Delta 9-tetrahydrocannabinol and cross-sensitization with morphine. *Psychopharmacology (Berl)* 2001; 158(3): 259-66.
- [80] Rubino T, Vigano D, Massi P, Parolaro D. The psychoactive ingredient of marijuana induces behavioural sensitization. *Eur J Neurosci* 2001; 14(5): 884-6.
- [81] Varvel SA, Martin BR, Lichtman AH. Lack of behavioral sensitization after repeated exposure to THC in mice and comparison to methamphetamine. *Psychopharmacology (Berl)* 2007; 193(4): 511-9.
- [82] Arnold JC, Topple AN, Hunt GE, McGregor IS. Effects of pre-exposure and co-administration of the cannabinoid receptor agonist CP 55,940 on behavioral sensitization to cocaine. *Eur J Pharmacol* 1998; 354(1): 9-16.
- [83] Ellgren M, Hurd YL, Franck J. Amphetamine effects on dopamine levels and behavior following cannabinoid exposure during adolescence. *Eur J Pharmacol* 2004; 497(2): 205-13.

- [84] Kolb B, Gorny G, Limebeer CL, Parker LA. Chronic treatment with Delta-9-tetrahydrocannabinol alters the structure of neurons in the nucleus accumbens shell and medial prefrontal cortex of rats. *Synapse* 2006; 60(6): 429-36.
- [85] Muschamp JW, Sivi SM. Behavioral sensitization to amphetamine follows chronic administration of the CB1 agonist WIN 55,212-2 in Lewis rats. *Pharmacol Biochem Behav* 2002; 73(4): 835-42.
- [86] Wiley JL, Evans RL, Grainger DB, Nicholson KL. Age-dependent differences in sensitivity and sensitization to cannabinoids and 'club drugs' in male adolescent and adult rats. *Addict Biol* 2008; 13(3-4): 277-86.
- [87] Cadoni C, Valentini V, Di Chiara G. Behavioral sensitization to delta 9-tetrahydrocannabinol and cross-sensitization with morphine: differential changes in accumbal shell and core dopamine transmission. *J Neurochem* 2008; 106(4): 1586-93.
- [88] Perrotti LI, Weaver RR, Robison B, *et al.* Distinct patterns of DeltaFosB induction in brain by drugs of abuse. *Synapse* 2008; 62(5): 358-69.
- [89] Fadda P, Robinson L, Fratta W, Pertwee RG, Riedel G. Differential effects of THC- or CBD-rich cannabis extracts on working memory in rats. *Neuropharmacology* 2004; 47(8): 1170-9.
- [90] Hayakawa K, Mishima K, Hazekawa M, *et al.* Cannabidiol potentiates pharmacological effects of Delta(9)-tetrahydrocannabinol via CB(1) receptor-dependent mechanism. *Brain Res* 2008; 1188: 157-64.
- [91] Heyser CJ, Hampson RE, Deadwyler SA. Effects of delta-9-tetrahydrocannabinol on delayed match to sample performance in rats: alterations in short-term memory associated with changes in task specific firing of hippocampal cells. *J Pharmacol Exp Ther* 1993; 264(1): 294-307.
- [92] Lichtman AH, Dimen KR, Martin BR. Systemic or intrahippocampal cannabinoid administration impairs spatial memory in rats. *Psychopharmacology (Berl)* 1995; 119(3): 282-90.
- [93] Campos AC, Guimaraes FS. Involvement of 5HT1A receptors in the anxiolytic-like effects of cannabidiol injected into the dorsolateral periaqueductal gray of rats. *Psychopharmacology (Berl)* 2008; 199(2): 223-30.
- [94] Campos AC, Guimaraes FS. Evidence for a potential role for TRPV1 receptors in the dorsolateral periaqueductal gray in the attenuation of the anxiolytic effects of cannabinoids. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33(8): 1517-21.
- [95] Deiana S, Watanabe A, Yamasaki Y, *et al.* Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarin (CBDV), Delta(9)-tetrahydrocannabinol (THC) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive-compulsive behaviour. *Psychopharmacology (Berl)* 2012; 219(3): 859-73.
- [96] Gomes FV, Resstel LB, Guimaraes FS. The anxiolytic-like effects of cannabidiol injected into the bed nucleus of the stria terminalis are mediated by 5-HT1A receptors. *Psychopharmacology (Berl)* 2011; 213(2-3): 465-73.
- [97] Guimaraes FS, Chiaretti TM, Graeff FG, Zuardi AW. Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology (Berl)* 1990; 100(4): 558-9.
- [98] Guimaraes FS, de Aguiar JC, Mechoulam R, Breuer A. Anxiolytic effect of cannabidiol derivatives in the elevated plus-maze. *Gen Pharmacol* 1994; 25(1): 161-4.
- [99] Moreira FA, Aguiar DC, Guimaraes FS. Anxiolytic-like effect of cannabidiol in the rat Vogel conflict test. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; 30(8): 1466-71.
- [100] Elbatsh MM, Assareh N, Marsden CA, Kendall DA. Anxiogenic-like effects of chronic cannabidiol administration in rats. *Psychopharmacology (Berl)* 2012.
- [101] Varvel SA, Wiley JL, Yang R, *et al.* Interactions between THC and cannabidiol in mouse models of cannabinoid activity. *Psychopharmacology (Berl)* 2006; 186(2): 226-34.
- [102] Guimaraes VM, Zuardi AW, Del Bel EA, Guimaraes FS. Cannabidiol increases Fos expression in the nucleus accumbens but not in the dorsal striatum. *Life Sci* 2004; 75(5): 633-8.
- [103] Malone DT, Hill MN, Rubino T. Adolescent cannabis use and psychosis: epidemiology and neurodevelopmental models. *Br J Pharmacol* 2010; 160: 511-22.
- [104] Leweke FM and Schneider M. Chronic pubertal cannabinoid treatment as a behavioural model for aspects of schizophrenia: effects of the atypical antipsychotic quetiapine. *Int J Neuropsychopharmacol* 2011; 14(1): 43-51.
- [105] Schneider M. Puberty as a highly vulnerable developmental period for the consequences of cannabis exposure. *Addict Biol* 2008; 13(2): 253-63.
- [106] Schneider M, Koch M. Chronic pubertal, but not adult chronic cannabinoid treatment impairs sensorimotor gating, recognition memory, and the performance in a progressive ratio task in adult rats. *Neuropsychopharmacology* 2003; 28(10): 1760-9.
- [107] Schneider M, Schomig E, Leweke FM. Acute and chronic cannabinoid treatment differentially affects recognition memory and social behavior in pubertal and adult rats. *Addict Biol* 2008; 13(3-4): 345-57.
- [108] O'Shea M, Singh ME, McGregor IS, Mallet PE. Chronic cannabinoid exposure produces lasting memory impairment and increased anxiety in adolescent but not adult rats. *J Psychopharmacol* 2004; 18(4): 502-8.
- [109] Quinn HR, Matsumoto I, Callaghan PD, *et al.* Adolescent rats find repeated Delta(9)-THC less aversive than adult rats but display greater residual cognitive deficits and changes in hippocampal protein expression following exposure. *Neuropsychopharmacology* 2008; 33(5): 1113-26.
- [110] Rubino T, Realini N, Braida D, *et al.* Changes in hippocampal morphology and neuroplasticity induced by adolescent THC treatment are associated with cognitive impairment in adulthood. *Hippocampus* 2009; 19(8): 763-72.
- [111] Schramm-Sapota NL, Cha YM, Chaudhry S, *et al.* Differential anxiogenic, aversive, and locomotor effects of THC in adolescent and adult rats. *Psychopharmacology (Berl)* 2007; 191(4): 867-77.
- [112] Dalton VS and Zavitsanou K. Cannabinoid effects on CB1 receptor density in the adolescent brain: an autoradiographic study using the synthetic cannabinoid HU210. *Synapse* 2010; 64(11): 845-54.
- [113] Verduran M, Dalton VS, Zavitsanou K. GABA(A) receptor density is altered by cannabinoid treatment in the hippocampus of adult but not adolescent rats. *Brain Res* 2010; 1351: 238-45.
- [114] Zavitsanou K, Wang H, Dalton VS, Nguyen V. Cannabinoid administration increases 5HT1A receptor binding and mRNA expression in the hippocampus of adult but not adolescent rats. *Neuroscience* 2010; 169(1): 315-24.
- [115] Dalton VS, Zavitsanou K. Differential treatment regimen-related effects of cannabinoids on D1 and D2 receptors in adolescent and adult rat brain. *J Chem Neuroanat* 2010; 40(4): 272-80.
- [116] Ellgren M, Artmann A, Tkalych O, *et al.* Dynamic changes of the endogenous cannabinoid and opioid mesocorticolimbic systems during adolescence: THC effects. *Eur Neuropsychopharmacol* 2008; 18(11): 826-834.
- [117] Altar CA, Jurata LW, Charles V, *et al.* Deficient hippocampal neuron expression of proteasome, ubiquitin, and mitochondrial genes in multiple schizophrenia cohorts. *Biol Psychiatry* 2005; 58(2): 85-96.
- [118] Gorriti MA, Ferrer B, del Arco I, *et al.* Acute delta9-tetrahydrocannabinol exposure facilitates quinpirole-induced hyperlocomotion. *Pharmacol Biochem Behav* 2005; 81(1): 71-7.
- [119] Gorriti MA, Rodriguez de Fonseca F, Navarro M, Palomo T. Chronic (-)-delta9-tetrahydrocannabinol treatment induces sensitization to the psychomotor effects of amphetamine in rats. *Eur J Pharmacol* 1999; 365(2-3): 133-42.
- [120] Lamarque S, Taghzouti K, Simon H. Chronic treatment with Delta(9)-tetrahydrocannabinol enhances the locomotor response to amphetamine and heroin. Implications for vulnerability to drug addiction. *Neuropsychopharmacology* 2001; 41(1): 118-29.
- [121] Carboni E, Imperato A, Perezzi L, Di Chiara G. Amphetamine, cocaine, phencyclidine and nomifensine increase extracellular dopamine concentrations preferentially in the nucleus accumbens of freely moving rats. *Neuroscience* 1989; 28(3): 653-61.
- [122] Carboni E, Spilewoy C, Vacca C, *et al.* Cocaine and amphetamine increase extracellular dopamine in the nucleus accumbens of mice lacking the dopamine transporter gene. *J Neurosci* 2001; 21(9): RC141: 1-4.
- [123] Moreira FA, Guimaraes FS. Cannabidiol inhibits the hyperlocomotion induced by psychotomimetic drugs in mice. *Eur J Pharmacol* 2005; 512(2-3): 199-205.
- [124] Zuardi AW, Rodrigues JA, Cunha JM. Effects of cannabidiol in animal models predictive of antipsychotic activity. *Psychopharmacology (Berl)* 1991; 104(2): 260-4.

- [125] Valvassori SS, Elias G, de Souza B, *et al.* Effects of cannabidiol on amphetamine-induced oxidative stress generation in an animal model of mania. *J Psychopharmacol* 2011; 25(2): 274-80.
- [126] Andreatza AC, Kauer-Sant'Anna M, Frey BN, *et al.* Effects of mood stabilizers on DNA damage in an animal model of mania. *J Psychiatry Neurosci* 2008; 33(6): 516-24.
- [127] Frey BN, Andreatza AC, Cereser KM, *et al.* Effects of mood stabilizers on hippocampus BDNF levels in an animal model of mania. *Life Sci* 2006; 79(3): 281-6.
- [128] Frey BN, Martins MR, Petronilho FC, *et al.* Increased oxidative stress after repeated amphetamine exposure: possible relevance as a model of mania. *Bipolar Disord* 2006; 8(3): 275-80.
- [129] Marek GJ, Behl B, Bessalov AY, *et al.* Glutamatergic (N-methyl-D-aspartate receptor) hypofrontality in schizophrenia: too little juice or a miswired brain? *Mol Pharmacol* 2010; 77(3): 317-26.
- [130] Cochran SM, Kennedy M, McKerchar CE, *et al.* Induction of metabolic hypofunction and neurochemical deficits after chronic intermittent exposure to phencyclidine: differential modulation by antipsychotic drugs. *Neuropsychopharmacology* 2003; 28(2): 265-75.
- [131] Egerton A, Reid L, McGregor S, *et al.* Subchronic and chronic PCP treatment produces temporally distinct deficits in attentional set shifting and prepulse inhibition in rats. *Psychopharmacology (Berl)* 2008; 198(1): 37-49.
- [132] Pratt JA, Winchester C, Egerton A, Cochran SM, Morris BJ. Modelling prefrontal cortex deficits in schizophrenia: implications for treatment. *Br J Pharmacol* 2008; 153 Suppl 1: S465-70.
- [133] Goonawardena AV, Robinson L, Hampson RE, Riedel G. Cannabinoid and cholinergic systems interact during performance of a short-term memory task in the rat. *Learn Mem* 2010; 17(10): 502-11.
- [134] Vigano D, Guidali C, Petrosino S, *et al.* Involvement of the endocannabinoid system in phencyclidine-induced cognitive deficits modelling schizophrenia. *Int J Neuropsychopharmacol* 2009; 12(5): 599-614.
- [135] Spano MS, Fadda P, Frau R, Fattore L, Fratta W. Cannabinoid self-administration attenuates PCP-induced schizophrenia-like symptoms in adult rats. *Eur Neuropsychopharmacol* 2010; 20(1): 25-36.
- [136] Long LE, Malone DT, Taylor DA. Cannabidiol Reverses MK-801-Induced Disruption of Prepulse Inhibition in Mice. *Neuropsychopharmacology* 2006; 31(4): 795-803.
- [137] Gururajan A, Taylor DA, Malone DT. Effect of cannabidiol in a MK-801-rodent model of aspects of schizophrenia. *Behav Brain Res* 2011; 222(2): 299-308.
- [138] Bendall S, Jackson HJ, Hulbert CA, McGorry PD. Childhood trauma and psychotic disorders: a systematic, critical review of the evidence. *Schizophr Bull* 2008; 34(3): 568-79.
- [139] van Winkel R, Stefanis NC, Myin-Germeys I. Psychosocial stress and psychosis. A review of the neurobiological mechanisms and the evidence for gene-stress interaction. *Schizophr Bull* 2008; 34(6): 1095-105.
- [140] Geyer MA and Ellenbroek B. Animal behavior models of the mechanisms underlying antipsychotic atypicality. *Prog Neuropsychopharmacol Biol Psychiatry* 2003; 27(7): 1071-9.
- [141] Geyer MA, Wilkinson LS, Humby T, Robbins TW. Isolation rearing of rats produces a deficit in prepulse inhibition of acoustic startle similar to that in schizophrenia. *Biol Psychiatry* 1993; 34(6): 361-72.
- [142] Varty GB, Powell SB, Lehmann-Masten V, Buell MR, Geyer MA. Isolation rearing of mice induces deficits in prepulse inhibition of the startle response. *Behav Brain Res* 2006; 169(1): 162-7.
- [143] Muchimapura S, Fulford AJ, Mason R, Marsden CA. Isolation rearing in the rat disrupts the hippocampal response to stress. *Neuroscience* 2002; 112(3): 697-705.
- [144] Muchimapura S, Mason R, Marsden CA. Effect of isolation rearing on pre- and post-synaptic serotonergic function in the rat dorsal hippocampus. *Synapse* 2003; 47(3): 209-17.
- [145] Malone DT, Kearn CS, Chongue L, Mackie K, Taylor DA. Effect of social isolation on CB1 and D2 receptor and fatty acid amide hydrolase expression in rats. *Neuroscience* 2008; 152(1): 265-72.
- [146] Robinson SA, Loiacono RE, Christopoulos A, Sexton PM, Malone DT. The effect of social isolation on rat brain expression of genes associated with endocannabinoid signaling. *Brain Res* 2010; 1343: 153-67.
- [147] Ellenbroek BA and Cools AR. Early maternal deprivation and prepulse inhibition: the role of the postdeprivation environment. *Pharmacol Biochem Behav* 2002; 73(1): 177-84.
- [148] Ellenbroek BA, de Bruin NM, van Den Kroonenburg PT, van Luijtelar EL, Cools AR. The effects of early maternal deprivation on auditory information processing in adult Wistar rats. *Biol Psychiatry* 2004; 55(7): 701-7.
- [149] Ellenbroek BA, Derks N, Park HJ. Early maternal deprivation retards neurodevelopment in Wistar rats. *Stress* 2005; 8(4): 247-57.
- [150] Ellenbroek BA, van den Kroonenburg PT, Cools AR. The effects of an early stressful life event on sensorimotor gating in adult rats. *Schizophr Res* 1998; 30(3): 251-60.
- [151] Roceri M, Hendriks W, Racagni G, Ellenbroek BA, Riva MA. Early maternal deprivation reduces the expression of BDNF and NMDA receptor subunits in rat hippocampus. *Mol Psychiatry* 2002; 7(6): 609-16.
- [152] Suarez J, Llorente R, Romero-Zerbo SY, *et al.* Early maternal deprivation induces gender-dependent changes on the expression of hippocampal CB(1) and CB(2) cannabinoid receptors of neonatal rats. *Hippocampus* 2009; 19(7): 623-32.
- [153] Suarez J, Rivera P, Llorente R, *et al.* Early maternal deprivation induces changes on the expression of 2-AG biosynthesis and degradation enzymes in neonatal rat hippocampus. *Brain Res* 2010; 1349: 162-73.
- [154] Lopez-Gallardo M, Lopez-Rodriguez AB, Llorente-Berzal A, *et al.* Maternal deprivation and adolescent cannabinoid exposure impact hippocampal astrocytes, CB1 receptors and brain-derived neurotrophic factor in a sexually dimorphic fashion. *Neuroscience* 2012; 204: 90-103.
- [155] Harley M, Kelleher I, Clarke M, *et al.* Cannabis use and childhood trauma interact additively to increase the risk of psychotic symptoms in adolescence. *Psychol Med* 2010; 40(10): 1627-34.
- [156] Llorente-Berzal A, Fuentes S, Gagliano H, *et al.* Sex-dependent effects of maternal deprivation and adolescent cannabinoid treatment on adult rat behaviour. *Addict Biol* 2011; 16(4): 624-37.
- [157] Llorente R, Arranz L, Marco EM, *et al.* Early maternal deprivation and neonatal single administration with a cannabinoid agonist induce long-term sex-dependent psychoimmunoenocrine effects in adolescent rats. *Psychoneuroendocrinology* 2007; 32(6): 636-50.
- [158] Hill MN and Gorzalka BB. Increased sensitivity to restraint stress and novelty-induced emotionality following long-term, high dose cannabinoid exposure. *Psychoneuroendocrinology* 2006; 31(4): 526-36.
- [159] Patel S, Cravatt BF, Hillard CJ. Synergistic interactions between cannabinoids and environmental stress in the activation of the central amygdala. *Neuropsychopharmacology* 2005; 30(3): 497-507.
- [160] Rodriguez de Fonseca F, Carrera MR, Navarro M, Koob GF, Weiss F. Activation of corticotropin-releasing factor in the limbic system during cannabinoid withdrawal. *Science* 1997; 276(5321): 2050-4.
- [161] Arnold JC, Dielenberg RA, McGregor IS. Cannabinoids increase conditioned ultrasonic vocalisations and cat odour avoidance in rats: strain differences in drug-induced anxiety. *Life Sci* 2010; 87(17-18): 572-8.
- [162] Lepore M, Liu X, Savage V, Matalon D, Gardner EL. Genetic differences in delta 9-tetrahydrocannabinol-induced facilitation of brain stimulation reward as measured by a rate-frequency curve-shift electrical brain stimulation paradigm in three different rat strains. *Life Sci* 1996; 58(25): PL365-72.
- [163] Arnold JC, Topple AN, Mallet PE, Hunt GE, McGregor IS. The distribution of cannabinoid-induced Fos expression in rat brain: differences between the Lewis and Wistar strain. *Brain Res* 2001; 921(1-2): 240-55.
- [164] Chesworth R, Downey L, Logge W, Killcross S, Karl T. Cognition in female transmembrane domain neuregulin 1 mutant mice. *Behav Brain Res* 2012; 226(1): 218-23.
- [165] Duffy L, Cappas E, Lai D, Boucher AA, Karl T. Cognition in transmembrane domain neuregulin 1 mutant mice. *Neuroscience* 2010; 170(3): 800-7.
- [166] Karl T, Burne TH, Van den Buuse M, Chesworth R. Do transmembrane domain neuregulin 1 mutant mice exhibit a reliable sensorimotor gating deficit? *Behav Brain Res* 2011; 223(2): 336-41.
- [167] Karl T, Duffy L, Scimone A, Harvey RP, Schofield PR. Altered motor activity, exploration and anxiety in heterozygous neuregulin

- 1 mutant mice: implications for understanding schizophrenia. *Genes Brain Behav* 2007; 6(7): 677-87.
- [168] Stefansson H, Sigurdsson E, Steinthorsdottir V, *et al.* Neuregulin 1 and susceptibility to schizophrenia. *Am J Hum Genet* 2002; 71(4): 877-92.
- [169] Corfas G, Roy K, Buxbaum JD. Neuregulin 1-erbB signaling and the molecular/cellular basis of schizophrenia. *Nat Neurosci* 2004; 7(6): 575-80.
- [170] Harrison PJ and Law AJ. Neuregulin 1 and schizophrenia: genetics, gene expression, and neurobiology. *Biol Psychiatry* 2006; 60(2): 132-40.
- [171] Mei L and Xiong WC. Neuregulin 1 in neural development, synaptic plasticity and schizophrenia. *Nat Rev Neurosci* 2008; 9(6): 437-52.
- [172] Long LE, Chesworth R, Arnold JC, Karl T. A follow-up study: acute behavioural effects of Delta(9)-THC in female heterozygous neuregulin 1 transmembrane domain mutant mice. *Psychopharmacology (Berl)* 2010; 211(3): 277-89.
- [173] Geyer MA, Krebs-Thomson K, Braff DL, Swerdlow NR. Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: a decade in review. *Psychopharmacology (Berl)* 2001; 156(2-3): 117-54.
- [174] Stadelmann AM, Roser P, Arning L, *et al.* Acute effects of delta9-tetrahydrocannabinol on the auditory evoked mismatch negativity are modulated by the NRG1 gene. *Pharmacopsychiatry* 2010; 43(5): 194-5.
- [175] Boucher AA, Hunt GE, Karl T, *et al.* Heterozygous neuregulin 1 mice display greater baseline and Delta(9)-tetrahydrocannabinol-induced c-Fos expression. *Neuroscience* 2007; 149(4): 861-70.
- [176] Sumner BE, Cruise LA, Slattery DA, *et al.* Testing the validity of c-fos expression profiling to aid the therapeutic classification of psychoactive drugs. *Psychopharmacology (Berl)* 2004; 171(3): 306-21.
- [177] Sheehan TP, Chambers RA, Russell DS. Regulation of affect by the lateral septum: implications for neuropsychiatry. *Brain Res Brain Res Rev* 2004; 46(1): 71-117.
- [178] Cheeta S, Kenny PJ, File SE. Hippocampal and septal injections of nicotine and 8-OH-DPAT distinguish among different animal tests of anxiety. *Prog Neuropsychopharmacol Biol Psychiatry* 2000; 24(7): 1053-67.
- [179] Cheeta S, Kenny PJ, File SE. The role of 5-HT1A receptors in mediating the anxiogenic effects of nicotine following lateral septal administration. *Eur J Neurosci* 2000; 12(10): 3797-802.
- [180] Nguyen QH and Wagner EJ. Estrogen differentially modulates the cannabinoid- induced presynaptic inhibition of amino acid neurotransmission in proopiomelanocortin neurons of the arcuate nucleus. *Neuroendocrinology* 2006; 84(2): 123-37.
- [181] Weickert CS, Miranda-Angulo AL, Wong J, *et al.* Variants in the estrogen receptor alpha gene and its mRNA contribute to risk for schizophrenia. *Hum Mol Genet* 2008; 17(15): 2293-309.
- [182] Wong J and Weickert CS. Transcriptional interaction of an estrogen receptor splice variant and ErbB4 suggests convergence in gene susceptibility pathways in schizophrenia. *J Biol Chem* 2009; 284(28): 18824-32.
- [183] Long LE, Chesworth R, Huang X, *et al.* Transmembrane domain Nrg1 mutant mice show altered susceptibility to the neurobehavioural actions of repeated THC exposure in adolescence. *Int J Neuropsychopharmacol*, 2012 Jan 9:1-13. [Epub ahead of print]
- [184] Abe Y, Namba H, Zheng Y, Nawa H. In situ hybridization reveals developmental regulation of ErbB1-4 mRNA expression in mouse midbrain: implication of ErbB receptors for dopaminergic neurons. *Neuroscience* 2009; 161(1): 95-110.
- [185] Caspi A, Moffitt TE, Cannon M, *et al.* Moderation of the effect of adolescent-onset cannabis use on adult psychosis by a functional polymorphism in the catechol-O-methyltransferase gene: longitudinal evidence of a gene X environment interaction. *Biol Psychiatry* 2005; 57(10): 1117-27.
- [186] O'Tuathaigh CM, Hryniewiecka M, Behan A, *et al.* Chronic adolescent exposure to Delta-9-tetrahydrocannabinol in COMT mutant mice: impact on psychosis-related and other phenotypes. *Neuropsychopharmacology* 2010; 35(11): 2262-73.
- [187] O'Tuathaigh CM, Clarke G, Walsh J, *et al.* Genetic vs. pharmacological inactivation of COMT influences cannabinoid-induced expression of schizophrenia-related phenotypes. *Int J Neuropsychopharmacol* 2011: 1-12.
- [188] Marquez B and Van Bambeke F. ABC multidrug transporters: target for modulation of drug pharmacokinetics and drug-drug interactions. *Curr Drug Targets* 2011; 12(5): 600-20.
- [189] Holland ML, Allen JD, Arnold JC. Interaction of plant cannabinoids with the multidrug transporter ABCC1 (MRP1). *Eur J Pharmacol* 2008; 591(1-3): 128-31.
- [190] Holland ML, Lau DT, Allen JD, Arnold JC. The multidrug transporter ABCG2 (BCRP) is inhibited by plant-derived cannabinoids. *Br J Pharmacol* 2007; 152(5): 815-24.
- [191] Holland ML, Panetta JA, Hoskins JM, *et al.* The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells. *Biochem Pharmacol* 2006; 71(8): 1146-54.
- [192] Zhu HJ, Wang JS, Markowitz JS, *et al.* Characterization of P-glycoprotein inhibition by major cannabinoids from marijuana. *J Pharmacol Exp Ther* 2006; 317(2): 850-7.
- [193] Doran A, Obach RS, Smith BJ, *et al.* The impact of P-glycoprotein on the disposition of drugs targeted for indications of the central nervous system: evaluation using the MDR1A/1B knockout mouse model. *Drug Metab Dispos* 2005; 33(1): 165-74.
- [194] Wang JS, Taylor R, Ruan Y, *et al.* Olanzapine penetration into brain is greater in transgenic Abcb1a P-glycoprotein-deficient mice than FVB1 (wild-type) animals. *Neuropsychopharmacology* 2004; 29(3): 551-7.
- [195] Bonhomme-Faivre L, Benyamina A, Reynaud M, Farinotti R, Abbara C. Disposition of Delta tetrahydrocannabinol in CF1 mice deficient in mdr1a P-glycoprotein. *Addict Biol* 2008; 13(3-4): 295-300.
- [196] Benyamina A, Bonhomme-Faivre L, Picard V, *et al.* Association between ABCB1 C3435T polymorphism and increased risk of cannabis dependence. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33(7): 1270-4.
- [197] Spiro AS, Wong A, Boucher AA, Arnold JC. Enhanced Brain Disposition and Effects of Δ^9 -Tetrahydrocannabinol in P-glycoprotein and Breast Cancer Resistance Protein Knockout Mice. *PLoS ONE*, 2012; 7(4): e35937
- [198] Long LE, Chesworth R, Huang X, *et al.* Distinct Neurobehavioural Effects of Cannabidiol in Transmembrane Domain Neuregulin 1 Mutant Mice. *PLoS ONE* 2012; 7(4): e34129.
- [199] Ballmaier M, Bortolato M, Rizzetti C, *et al.* Cannabinoid receptor antagonists counteract sensorimotor gating deficits in the phencyclidine model of psychosis. *Neuropsychopharmacology* 2007; 32(10): 2098-107.
- [200] Martin RS, Secchi RL, Sung E, *et al.* Effects of cannabinoid receptor ligands on psychosis-relevant behavior models in the rat. *Psychopharmacology (Berl)* 2003; 165(2): 128-35.
- [201] Black MD, Stevens RJ, Rogacki N, *et al.* AVE1625, a cannabinoid CB1 receptor antagonist, as a co-treatment with antipsychotics for schizophrenia: improvement in cognitive function and reduction of antipsychotic-side effects in rodents. *Psychopharmacology (Berl)* 2011; 215(1): 149-63.
- [202] Guidali C, Vigano D, Petrosino S, *et al.* Cannabinoid CB1 receptor antagonism prevents neurochemical and behavioural deficits induced by chronic phencyclidine. *Int J Neuropsychopharmacol* 2011; 14(1): 17-28.
- [203] Chiang YC, Chen JC. The role of the cannabinoid type 1 receptor and down-stream cAMP/DARPP-32 signal in the nucleus accumbens of methamphetamine-sensitized rats. *J Neurochem* 2007; 103(6): 2505-17.
- [204] Corbille AG, Valjent E, Marsicano G, *et al.* Role of cannabinoid type 1 receptors in locomotor activity and striatal signaling in response to psychostimulants. *J Neurosci* 2007; 27(26): 6937-47.
- [205] Landa L, Sulcova A, Slais K. Involvement of cannabinoid CB1 and CB2 receptor activity in the development of behavioural sensitization to methamphetamine effects in mice. *Neuro Endocrinol Lett* 2006; 27(1-2): 63-9.
- [206] Thiemann G, Di Marzo V, Molleman A, Hasenohrl RU. The CB(1) cannabinoid receptor antagonist AM251 attenuates amphetamine-induced behavioural sensitization while causing monoamine changes in nucleus accumbens and hippocampus. *Pharmacol Biochem Behav* 2008; 89(3): 384-91.
- [207] Vlachou S, Nomikos GG, Panagis G. CB1 cannabinoid receptor agonists increase intracranial self-stimulation thresholds in the rat. *Psychopharmacology (Berl)* 2005; 179(2): 498-508.
- [208] Eisenstein SA, Holmes PV, Hohmann AG. Endocannabinoid modulation of amphetamine sensitization is disrupted in a rodent

- model of lesion-induced dopamine dysregulation. *Synapse* 2009; 63(11): 941-50.
- [209] Cador M, Bijiou Y, Stinus L. Evidence of a complete independence of the neurobiological substrates for the induction and expression of behavioral sensitization to amphetamine. *Neuroscience* 1995; 65(2): 385-95.
- [210] Vanderschuren LJ and Kalivas PW. Alterations in dopaminergic and glutamatergic transmission in the induction and expression of behavioral sensitization: a critical review of preclinical studies. *Psychopharmacology (Berl)* 2000; 151(2-3): 99-120.
- [211] Vezina P and Stewart J. Amphetamine administered to the ventral tegmental area but not to the nucleus accumbens sensitizes rats to systemic morphine: lack of conditioned effects. *Brain Res* 1990; 516(1): 99-106.
- [212] Ramiro-Fuentes S and Fernandez-Espejo E. Sensitization to cocaine is inhibited after intra-accumbal GR103691 or rimonabant, but it is enhanced after co-infusion indicating functional interaction between accumbens D(3) and CB1 receptors. *Psychopharmacology (Berl)* 2011; 214(4): 949-59.
- [213] Chavarria-Siles I, Contreras-Rojas J, Hare E, *et al.* Cannabinoid receptor 1 gene (CNR1) and susceptibility to a quantitative phenotype for hebephrenic schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 2008; 147(3): 279-84.
- [214] Leroy S, Griffon N, Bourdel MC, *et al.* Schizophrenia and the cannabinoid receptor type 1 (CB1): association study using a single-base polymorphism in coding exon 1. *Am J Med Genet* 2001; 105(8): 749-52.
- [215] Martinez-Gras I, Hoenicka J, Ponce G, *et al.* (AAT)n repeat in the cannabinoid receptor gene, CNR1: association with schizophrenia in a Spanish population. *Eur Arch Psychiatry Clin Neurosci* 2006; 256(7): 437-41.
- [216] Ujike H, Takaki M, Nakata K, *et al.* CNR1, central cannabinoid receptor gene, associated with susceptibility to hebephrenic schizophrenia. *Mol Psychiatry* 2002; 7(5): 515-8.
- [217] Seifert J, Ossege S, Emrich HM, Schneider U, Stuhmann M. No association of CNR1 gene variations with susceptibility to schizophrenia. *Neurosci Lett* 2007; 426(1): 29-33.
- [218] Tsai SJ, Wang YC, Hong CJ. Association study of a cannabinoid receptor gene (CNR1) polymorphism and schizophrenia. *Psychiatr Genet* 2000; 10(3): 149-51.
- [219] Zammit S, Spurlock G, Williams H, *et al.* Genotype effects of CHRNA7, CNR1 and COMT in schizophrenia: interactions with tobacco and cannabis use. *Br J Psychiatry* 2007; 191: 402-7.
- [220] Dalton VS, Long LE, Weickert CS, Zavitsanou K. Paranoid schizophrenia is characterized by increased CB1 receptor binding in the dorsolateral prefrontal cortex. *Neuropsychopharmacology* 2011; 36(8): 1620-30.
- [221] Dean B, Sundram S, Bradbury R, Scarr E, Copolov D. Studies on [3H]CP-55940 binding in the human central nervous system: regional specific changes in density of cannabinoid-1 receptors associated with schizophrenia and cannabis use. *Neuroscience* 2001; 103(1): 9-15.
- [222] Newell KA, Deng C, Huang XF. Increased cannabinoid receptor density in the posterior cingulate cortex in schizophrenia. *Exp Brain Res* 2006; 172(4): 556-60.
- [223] Zavitsanou K, Garrick T, Huang XF. Selective antagonist [3H]SR141716A binding to cannabinoid CB1 receptors is increased in the anterior cingulate cortex in schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2004; 28(2): 355-60.
- [224] Ledent C, Valverde O, Cossu G, *et al.* Unresponsiveness to cannabinoids and reduced addictive effects of opiates in CB1 receptor knockout mice. *Science* 1999; 283(5400): 401-4.
- [225] Zimmer A, Zimmer AM, Hohmann AG, Herkenham M, Bonner TI. Increased mortality, hypoactivity, and hypoalgesia in cannabinoid CB1 receptor knockout mice. *Proc Natl Acad Sci U S A* 1999; 96(10): 5780-5.
- [226] Reibaud M, Obinu MC, Ledent C, *et al.* Enhancement of memory in cannabinoid CB1 receptor knock-out mice. *Eur J Pharmacol* 1999; 379(1): R1-2.
- [227] Varvel SA, Lichtman AH. Evaluation of CB1 receptor knockout mice in the Morris water maze. *J Pharmacol Exp Ther* 2002; 301(3): 915-24.
- [228] Varvel SA, Anum EA, Lichtman AH. Disruption of CB(1) receptor signaling impairs extinction of spatial memory in mice. *Psychopharmacology (Berl)* 2005; 179(4): 863-72.
- [229] Haller J, Bakos N, Szirmay M, Ledent C, Freund TF. The effects of genetic and pharmacological blockade of the CB1 cannabinoid receptor on anxiety. *Eur J Neurosci* 2002; 16(7): 1395-8.
- [230] Haller J, Szirmai M, Varga B, Ledent C, Freund TF. Cannabinoid CB1 receptor dependent effects of the NMDA antagonist phencyclidine in the social withdrawal model of schizophrenia. *Behav Pharmacol* 2005; 16(5-6): 415-22.
- [231] Haller J, Varga B, Ledent C, Freund TF. CB1 cannabinoid receptors mediate anxiolytic effects: convergent genetic and pharmacological evidence with CB1-specific agents. *Behav Pharmacol* 2004; 15(4): 299-304.
- [232] Ferrer B, Gorriti MA, Palomino A, *et al.* Cannabinoid CB1 receptor antagonism markedly increases dopamine receptor-mediated stereotypies. *Eur J Pharmacol* 2007; 559(2-3): 180-3.
- [233] Mato S, Aso E, Castro E, *et al.* CB1 knockout mice display impaired functionality of 5-HT1A and 5-HT2A/C receptors. *J Neurochem* 2007; 103(5): 2111-20.
- [234] Ishiguro H, Horiuchi Y, Ishikawa M, *et al.* Brain Cannabinoid CB2 Receptor in Schizophrenia. *Biological Psychiatry* 2010; 67(10): 974-82.
- [235] Minocci D, Massey J, Martino A, *et al.* Genetic association between bipolar disorder and 524A>C (Leu133Ile) polymorphism of CNR2 gene, encoding for CB2 cannabinoid receptor. *J Affect Disord* 2011; 134(1-3): 427-30.
- [236] Ortega-Alvaro A, Aracil-Fernandez A, Garcia-Gutierrez MS, Navarrete F, Manzanares J. Deletion of CB2 cannabinoid receptor induces schizophrenia-related behaviors in mice. *Neuropsychopharmacology* 2011; 36(7): 1489-504.
- [237] ElSohly MA, Ross SA, Mehmedic Z, *et al.* Potency trends of delta9-THC and other cannabinoids in confiscated marijuana from 1980-1997. *J Forensic Sci* 2000; 45(1): 24-30.
- [238] Mehmedic Z, Chandra S, Slade D, *et al.* Potency trends of Delta9-THC and other cannabinoids in confiscated cannabis preparations from 1993 to 2008. *J Forensic Sci* 2010; 55(5): 1209-17.
- [239] Pijlman FT, Rigter SM, Hoek J, Goldschmidt HM, Niesink RJ. Strong increase in total delta-THC in cannabis preparations sold in Dutch coffee shops. *Addict Biol* 2005; 10(2): 171-80.
- [240] Potter DJ, Clark P, Brown MB. Potency of delta 9-THC and other cannabinoids in cannabis in England in 2005: implications for psychoactivity and pharmacology. *J Forensic Sci* 2008; 53(1): 90-4.
- [241] Vann RE, Gamage TF, Warner JA, *et al.* Divergent effects of cannabidiol on the discriminative stimulus and place conditioning effects of Delta(9)-tetrahydrocannabinol. *Drug Alcohol Depend* 2008; 94(1-3): 191-8.
- [242] Zuardi AW, Karniol IG. Effects on variable-interval performance in rats of delta 9-tetrahydrocannabinol and cannabidiol, separately and in combination. *Braz J Med Biol Res* 1983; 16(2): 141-6.
- [243] Zuardi AW, Teixeira NA, Karniol IC. Pharmacological interaction of the effects of delta 9-trans-tetrahydrocannabinol and cannabidiol on serum corticosterone levels in rats. *Arch Int Pharmacodyn Ther* 1984; 269(1): 12-9.
- [244] Fernandes M, Schabarek A, Coper H, Hill R. Modification of delta9-THC-actions by cannabinal and cannabidiol in the rat. *Psychopharmacologia* 1974; 38(4): 329-38.
- [245] Hiltunen AJ, Jarbe TU. Interactions between delta 9-tetrahydrocannabinol and cannabidiol as evaluated by drug discrimination procedures in rats and pigeons. *Neuropharmacology* 1986; 25(2): 133-42.
- [246] Klein C, Karanges E, Spiro A, *et al.* Cannabidiol potentiates Delta-tetrahydrocannabinol (THC) behavioural effects and alters THC pharmacokinetics during acute and chronic treatment in adolescent rats. *Psychopharmacology (Berl)* 2011; 218(2): 443-57.
- [247] Finn DP, Beckett SR, Roe CH, *et al.* Effects of coadministration of cannabinoids and morphine on nociceptive behaviour, brain monoamines and HPA axis activity in a rat model of persistent pain. *Eur J Neurosci* 2004; 19(3): 678-86.
- [248] Morgan CJ, Schafer G, Freeman TP, Curran HV. Impact of cannabidiol on the acute memory and psychotomimetic effects of smoked cannabis: naturalistic study: naturalistic study [corrected]. *Br J Psychiatry* 2010; 197(4): 285-90.
- [249] Wong A, Gunasekaran N, Hancock DP, *et al.* The major plant-derived cannabinoid Δ9-tetrahydrocannabinol promotes hypertrophy and macrophage infiltration in adipose tissue. *Horm Metab Res* 2012; 44: 105-113.

- [250] Russo EB, Burnett A, Hall B, Parker KK. Agonistic properties of cannabidiol at 5-HT_{1a} receptors. *Neurochem Res* 2005; 30(8): 1037-43.
- [251] Bornheim LM, Kim KY, Li J, Perotti BY, Benet LZ. Effect of cannabidiol pretreatment on the kinetics of tetrahydrocannabinol metabolites in mouse brain. *Drug Metab Dispos* 1995; 23(8): 825-31.
- [252] Jones G, Pertwee RG. A metabolic interaction *in vivo* between cannabidiol and 1 -tetrahydrocannabinol. *Br J Pharmacol* 1972; 45(2): 375-7.
- [253] Reid MJ and Bornheim LM. Cannabinoid-induced alterations in brain disposition of drugs of abuse. *Biochem Pharmacol* 2001; 61(11): 1357-67.
- [254] Bornheim LM and Correia MA. Selective inactivation of mouse liver cytochrome P-450_{III}A by cannabidiol. *Mol Pharmacol* 1990; 38(3): 319-26.
- [255] Bornheim LM, Everhart ET, Li J, Correia MA. Characterization of cannabidiol-mediated cytochrome P450 inactivation. *Biochem Pharmacol* 1993; 45(6): 1323-31.
- [256] Bornheim LM and Grillo MP. Characterization of cytochrome P450 3A inactivation by cannabidiol: possible involvement of cannabidiol-hydroxyquinone as a P450 inactivator. *Chem Res Toxicol* 1998; 11(10): 1209-16.

Received: March 30, 2012

Accepted: April 18, 2012