

# Differences between Smokers and Nonsmokers in Regional Gray Matter Volumes and Densities

Arthur L. Brody, Mark A. Mandelkern, Murray E. Jarvik, Grace S. Lee, Eryln C. Smith, Joe C. Huang, Robert G. Bota, George Bartzokis, and Edythe D. London

**Background:** Magnetic resonance imaging (MRI) studies have demonstrated large-scale brain abnormalities in cigarette smokers, such as ventricular enlargement and atrophy. Converging lines of evidence point to functional differences between smokers and nonsmokers in specific brain regions, namely the lateral prefrontal cortex (PFC), anterior cingulate cortex (ACC), ventral striatum, and thalamus. Using MRI, we examined these regions for differences in gray matter between smokers and nonsmokers.

**Methods:** Thirty-six otherwise healthy adults (19 smokers and 17 nonsmoking control subjects) underwent three-dimensional Fourier-transform spoiled-gradient-recalled acquisition MRI of the brain. Both hand-drawn regions of interest and the computer program voxel-based morphometry were used to assess group differences in regional gray matter volumes and densities, respectively.

**Results:** Smokers had smaller gray matter volumes and lower gray matter densities than nonsmokers in the PFC bilaterally, along with smaller volumes in the left dorsal ACC and lower gray matter densities in the right cerebellum. Smokers also had negative associations between pack-year smoking history and PFC gray matter densities.

**Conclusions:** Smokers and nonsmokers differed in regional gray matter in brain areas previously linked with nicotine dependence. These findings might reflect effects of chronic smoking, predisposing traits that lead to smoking, or some combination of these factors. Biol Psychiatry 2004;55:77–84 © 2004 Society of Biological Psychiatry

**Key Words:** Magnetic resonance imaging, nicotine dependence, voxel-based morphometry, prefrontal cortex, anterior cingulate cortex, ventral striatum

Magnetic resonance imaging (MRI) studies have demonstrated that cigarette smoking history is associated with large-scale structural brain abnormalities. Cigarette usage, as indexed by pack-year smoking history, has been linked with increased sulcal and ventricular grade (Longstreth et al 2000) and atrophy (Longstreth et al 2001) in the elderly. Smoking history is also associated with severity of periventricular white matter hyperintensities in most (Fukuda and Kitani 1996; Kobayashi et al 1997; Liao et al 1997; Longstreth et al 2000; Tsushima et al 2002) but not all (Yamashita et al 1996; Yetkin et al 1993) studies in which this question was examined. In addition, smoking is a risk factor for both silent (Howard et al 1998; Longstreth et al 1998) and symptomatic (Longstreth et al 1998, 2001) stroke. Although these studies have examined large-scale abnormalities, no one has yet reported regional gray matter volume and density differences between otherwise healthy smokers and nonsmoking control subjects.

Converging lines of evidence point to four candidate brain regions that might differ in gray matter volumes and densities between smokers and nonsmokers. Functional MRI (fMRI) and positron emission tomography (PET) studies demonstrate that administration of nicotine (or cigarette smoking) increases activity in the prefrontal cortex (PFC) (Nakamura et al 2000; Stein et al 1998), anterior cingulate cortex (ACC) (Stein et al 1998), ventral striatum (Nakamura et al 2000; Stein et al 1998), and

thalamus (Domino et al 2000; Nakamura et al 2000; Rose et al 2003; Stein et al 1998; Zubieta et al 2001). Functional imaging studies also show that smokers exposed to cigarette-related cues have activation of the PFC (Due et al 2002) and ACC (Brody et al 2002). Consistent with these findings in humans are imaging studies of nonhuman primates and rodents, which demonstrate that nicotine administration results in dopamine release in the ventral striatum (Dewey et al 1999; Di Chiara and Imperato 1988) and that the highest density of nicotinic acetylcholine receptors is in the thalamus (Clarke et al 1984; Dávila-García et al 1997; Horti et al 1998; London et al 1985, 1995; Musachio et al 1997). Thus, four regions (the PFC, ACC, ventral striatum, and thalamus) are implicated repeatedly in responses to smoking and smoking cues.

In addition to direct effects of smoking and smoking cues on the candidate brain regions, indirect evidence from behavioral studies strengthens the association between these regions and smoking status. Smokers do not perform as well as nonsmokers on tasks that require working memory (N-back type) (Ernst et al 2001; Spilich et al 1992), which are known to activate the lateral PFC (Braver et al 2001; Meyer-Lindenberg et al 2001; Zurovski et al 2002). Smokers also have higher rates than the general population of impulsivity (Mitchell 1999; Waldeck and Miller 1997), a trait associated with diminished PFC function (Brower and Price 2001). Additionally, smokers have state-dependent (abstinence vs. satiety) alterations in reaction time (Hatsukami et al 1989; Pritchard et al 1992; Shiffman et al 1995), arousal (Parrott and Kaye 1999), motivation (Powell et al 2002), and sustained attention (Rusted et al 2000), which are all functions of the ACC (Bench et al 1993; Critchley et al 2001; Rauch et al 1999).

We are not aware of any previous reports focusing on regional gray matter in smokers; however, associations between regional gray matter and demographic (and other substance dependence) variables have been reported. Prior studies have examined associations between regional brain volumes and age (Bartzokis et al 2001; Coffey et al 1992; Cowell et al 1994; Jernigan et al 1991, 2001; Mueller et al 1998; Murphy et al 1996; Pfefferbaum et al 1998; Raz et al 1997; Salat et al 2001; Tisserand et al 2000; Van Der Werf et al 2001), gender (Cowell et al 1994; Gur et al 2002a, 2002b; Murphy et al 1996), and intelligence

From the Veterans Affairs Greater Los Angeles Healthcare System (ALB, MAM, MEJ, GSL, ECS, JCH, RGB, GB, EDL); Department of Psychiatry and Biobehavioral Sciences (ALB, MEJ, EDL), Department of Neurology and Laboratory of Neuroimaging (GB), and Department of Molecular and Medical Pharmacology (EDL), University of California at Los Angeles, at Los Angeles; and Department of Physics (MAM), University of California at Irvine, Irvine, California.

Address reprint requests to Arthur L. Brody, M.D., 300 UCLA Medical Plaza, Suite 2200, Los Angeles CA 90095.

Received February 13, 2003; revised May 22, 2003; accepted June 6, 2003.

(Andreasen et al 1993; Reiss et al 1996), as well as other substance dependencies (Bartzokis et al 2000; Fein et al 2002b; Jacobsen et al 2001a, 2001b; Liu et al 1998). None of these studies reported the effects of smoking status in detail, although one of the aging studies did use smoking status (along with other cardiovascular risk factors) as a single covariate in the statistical analysis (Coffey et al 1992).

## Methods and Materials

### Subjects

Thirty-six subjects (19 smokers and 17 nonsmoking control subjects), 21–65 years of age, were recruited through local newspaper advertisements. They were initially screened during a semistructured telephone interview to assess smoking, medical, psychiatric, medication, and substance use history. Smokers were defined as those who smoked at least 20 cigarettes per day and met DSM-IV criteria for nicotine dependence. Nonsmoking history was defined as having smoked no more than five cigarettes in a lifetime. Subjects were also screened on the telephone for exclusion criteria (see below). Those who passed these initial screening criteria met with the principal investigator (ALB), and all subjects who participated in the study provided written informed consent after the procedures had been fully explained. The consent form and study procedures were approved by the institutional review board for the Greater Los Angeles Veterans Affairs Healthcare System. Subjects in this study also participated in a separate PET imaging study (Brody et al 2002).

Subjects were excluded for any lifetime history of medical, psychiatric, substance abuse, or medication use thought to affect brain structure or function at the time of scanning. Examples of medical exclusion criteria include primarily neurologic conditions, such as a history of epilepsy or seizures, stroke, or head trauma with loss of consciousness. Potential research participants with any lifetime history of an Axis I psychiatric disorder, as diagnosed by screening questions from the Structured Clinical Interview for DSM-IV (First et al 1995), were excluded. Substance abuse exclusion criteria included alcohol use more than 1 drink equivalent per day, any illicit drug use other than marijuana, or marijuana use more than once per month. Urine toxicology screens were obtained on all subjects with questionable drug abuse history, and subjects were excluded from further participation if their urine toxicology screen was positive. Medication exclusions included any history of regular psychotropic drug usage or any current regular usage of medications with the potential to affect the central nervous system (e.g.,  $\beta$ -blockers or analgesic medications).

Clinical ratings obtained before scanning included the 17-item Hamilton Depression Rating Scale (HAM-D; Hamilton 1967), the Hamilton Anxiety Rating Scale (HAM-A; Hamilton 1969), and the Fagerström Test for Nicotine Dependence (FTND; Fagerström 1978; Heatherton et al 1991). Exhaled carbon monoxide levels (Bedfont EC-50, Microsmokerlyzer II, Kent, United Kingdom) were obtained as a rough measure of recent cigarette smoking at the time that rating scales were administered. Handedness was determined with a standard rating scale (Oldfield 1971).

### MRI

An MRI scan of the brain was obtained for each subject within 3 days of the clinical interview. The scans were performed on a Vision 1.5-Tesla (Siemens Medical Systems, Iselin, New Jersey)

scanner with the following protocol: three-dimensional Fourier-transform spoiled-gradient-recalled acquisition, repetition time = 30 msec, echo time = 7 msec, 30° angle, 256 × 192 view matrix, 23 cm field of view. The acquired volume was reconstructed as contiguous 1.5-mm-thick transaxial slices. Subjects were instructed not to smoke on the morning of the scan; all scans were performed at 7:00 AM, with subjects arriving at the scanner at 6:30 AM.


### MRI Scan Analysis

MRI data were analyzed with two complementary methods: 1) hand-drawn regions of interest (ROIs) to determine gray matter volumes of specific brain structures; and 2) voxel-based morphometry (VBM) (Ashburner and Friston 2000) to measure relative gray matter density. Each method has strengths and limitations (Ashburner and Friston 2001; Bookstein 2001; Friston et al 1994; Steinmetz and Seitz 1991; Tisserand et al 2002). The ROI method gives direct measurement of gray matter volumes of selected structures but is limited by the subjectivity of investigator-defined regional boundaries and the potential for errors in region drawing. Volume-based morphometry measures tissue composition (e.g., gray matter density) and has the advantage of surveying the entire brain without the use of manual techniques; however, this method relies on spatial normalization to a standard brain template, thus deforming structures of interest and comingling effects of tissue composition with those of regional boundaries. Volume-based morphometry has been used to determine differences between cocaine-dependent subjects and normal control subjects (Franklin et al 2002) and has been validated against ROI methods (Good et al 2002; Karas et al 2002) and known structural abnormalities in subjects with degenerative dementias (Burton et al 2002; Busatto et al 2003).

### Regions of Interest

For the ROI analysis, six ROIs were hand drawn bilaterally on transaxial MRI slices (Figure 1) by a team of investigators blind to subject identity (GSL, ECS, JCH, and RGN), based on a priori hypotheses derived from the literature cited above. The ROIs were the dorsolateral and ventrolateral prefrontal cortices (DLPFC and VLPFC), dorsal and ventral ACC, ventral striatum, and thalamus. Boundaries for these regions were defined according to standard atlases (Damasio 1995; Mai et al 1997). Because subtle differences in regional boundaries can alter results, each ROI was assigned to a specific region drawer, and all ROIs were reviewed twice by the region drawers and the principal investigator (blinded to subject identity) to improve uniformity of placement. Intrarater reliability with this method (based on replicating three complete ROI sets) was .95.

The DLPFC (roughly corresponding to Brodmann areas [BA] 46/9) and VLPFC (roughly corresponding to BA 45/10) were drawn as the dorsal and ventral halves of the middle frontal gyrus (gray matter), respectively, and were drawn in approximately 10 planes each. The border between DLPFC and VLPFC was at the level of the middle of the body of the caudate nucleus. The anterior border of DLPFC and VLPFC was the superior frontal sulcus, and the posterior border was the inferior frontal sulcus. The inferior border of VLPFC was the level directly above the eyes, where the frontal cortex takes on a convex form unlike its concave appearance in the lower planes. The superior border of the DLPFC was at the level of the middle of the body of the cingulate gyrus. The ventral striatum was drawn in eight planes



**Figure 1.** Regions of interest drawn on magnetic resonance images of a study subject. DLPFC, dorsolateral prefrontal cortex; VLPFC, ventrolateral prefrontal cortex; Dorsal ACC, dorsal anterior cingulate cortex; Ventral ACC, ventral anterior cingulate cortex; Striat, striatum; Thal, thalamus.

and consisted of the ventral half of the head of caudate nucleus/nucleus accumbens. The entire thalamus was drawn in 12 planes. Supratentorial whole brain was also drawn in approximately 40 planes so that ratios of ROI to global volume could be determined.

For each ROI, volume was determined by multiplying the area of each drawn region ( $\text{mm}^2$ ) by the MRI slice thickness (1.5 mm) and summing the volumes of all slices. Region of interest volumes were then divided by whole brain volumes to determine the ratio ROI/global (normalized ROI value) for use in statistical analysis. The distribution and variance of all normalized ROI values were examined and determined to be suitable for parametric analyses.

### Voxel-Based Morphometry

The standard method of VBM was performed to determine group differences in regional gray matter density (Ashburner and Friston 2000). To prepare images for VBM, skull stripping was first performed with the use of an automated algorithm from the MEDx 3.3 software program (Sensor Systems, Sterling, Virginia). Statistical parametric mapping (SPM99) was then used to spatially normalize all images to a standardized template obtained from the developers of SPM. Images were then partitioned into gray matter, white matter, and cerebro-

spinal fluid, with the gray matter partitions being used for analysis (Ashburner and Friston 1997). The SPM option for inhomogeneity correction was applied, along with a 12-mm full width half maximum smoothing kernel and proportional scaling for statistical analysis.

### Statistical Analyses

Means ( $\pm$  SDs) were determined for clinical variables of interest and for normalized ROI values. Percent differences in normalized ROI value between smokers and nonsmokers were calculated as the difference between normalized ROI values for the groups divided by the mean normalized ROI value of the groups  $\times$  100.

For the ROI analysis, an overall multivariate analysis of covariance (MANCOVA) was performed for the six regions of interest bilaterally, with normalized ROI values used as the dependent variables, smoking status (smoker vs. nonsmoker) as a between-subjects factor, and age and gender as covariates (because of the known effects of these variables on regional brain volumes, as cited above). Based on the MANCOVA results, univariate ANCOVAs were performed for individual ROIs, with the same between-subjects factor and covariates as above, to determine which regions accounted

for the overall significant group difference. Alpha levels were set at  $p < .05$  (two-tailed).

A statistical analysis similar to the ROI analysis was performed with VBM (using SPM99) to explore the whole brain for differences in regional gray matter density between smokers and nonsmoking control subjects. The “compare groups” SPM option was used, with age and gender as nuisance covariates, to directly determine regions in which smokers and nonsmokers differed in gray matter density. A statistical threshold of  $p < .001$ , uncorrected for multiple comparisons, was used for voxels of peak significance, along with a minimum extent threshold of 50 contiguous voxels.

## Results

The smoker and nonsmoker groups were similar in age (mean  $39.5 \pm 10.3$  vs.  $37.9 \pm 12.9$  years, respectively), gender (42.1% vs. 41.2% female), ethnicity (74% vs. 64% Caucasian), handedness (0% vs. 6% left-handed), and HAM-D (mean  $2.0 \pm 2.7$  vs.  $1.2 \pm 1.7$ ) and HAM-A ( $2.4 \pm 2.3$  vs.  $1.6 \pm 1.8$ ) scores. Smokers had higher exhaled carbon monoxide levels ( $18.3 \pm 7.1$  vs.  $1.9 \pm .7$  parts per million (ppm), Student  $t$  test,  $p = 7.1 \times 10^{-9}$ ), and FTND scores ( $5.1 \pm 1.9$  vs.  $0 \pm 0$ , Student  $t$  test,  $p = 6.5 \times 10^{-13}$ ) than nonsmokers. Smokers smoked an average of  $26.2 \pm 7.4$  cigarettes per day (range, 20–40) and had a mean  $31.0 \pm 17.9$  pack-year smoking history (range, 9–70).

In the overall MANCOVA in which normalized ROIs were used, a significant effect of smoking status was found [ $F(12,21) = 3.0$ ,  $p = .013$ ]. Analyses of covariance performed for individual ROIs revealed that the left DLPFC [ $F(1,32) = 6.1$ ,  $p = .019$ ], left and right VLPFC [ $F(1,32) = 8.1$ ,  $p = .008$  and  $F(1,32) = 8.9$ ,  $p = .005$ , respectively], and left dorsal ACC [ $F(1,32) = 4.6$ ,  $p = .039$ ] had significant between-group differences, indicating that smokers had smaller PFC and left dorsal ACC gray matter volumes than nonsmokers (Table 1). For the DLPFC, smokers had a mean 15.7% smaller normalized gray matter volume on the left and 11.9% on the right. For the VLPFC, smokers had a mean 19.2% smaller gray matter volume on the left and 21.2% on the right. The left dorsal ACC was 18.5% smaller in smokers than nonsmokers. The remaining regions did not have significant between-group differences.

In the VBM analysis, gray matter density was lower in smokers than in nonsmokers in the prefrontal cortices (primarily the bilateral DLPFC but also the left VLPFC) (Table 2 and Figure 2) and also the right medial cerebellum. Not only did a group of voxel clusters pass the predetermined threshold for significance, but clusters in both the left and right DLPFC contained voxels that were significant after the Bonferroni-type correction within the SPM program ( $Z = 6.66$ ,  $x, y, z = -32, 40, 36$ ,  $p = .006$ ; and  $Z = 6.19$ ,  $x, y, z = 24, 42, 32$ ,  $p = .017$ , corrected for multiple comparisons, for the left and right, respectively). There were no regions of gray matter density significantly higher in smokers than in nonsmokers.

As a follow-up to these central study findings, we performed a preliminary analysis to determine associations between pack-year smoking history and regional gray matter volume and density for regions found to be significant in the preceding analyses, while recognizing that this analysis would be limited by a smaller sample size. For the four ROIs found to have significant between-group differences in the above analysis, none had significant Pearson correlation coefficients with pack-year smoking history (all  $p$  values  $\geq .46$ ). For the VBM analysis of main effect of pack-year smoking history on regional gray matter

**Table 1.** Comparison of Normalized Regions of Interest Values (ROI/Global) between Smokers and Nonsmokers

Region of Interest	Smokers (Mean $\pm$ SD)	Nonsmokers (Mean $\pm$ SD)	F Value (Smoking Status)
DLPFC			
L	9.2 ( $\pm$ 1.6)	10.7 ( $\pm$ 2.1)	6.1 <sup>a</sup>
R	9.4 ( $\pm$ 1.8)	10.5 ( $\pm$ 1.8)	3.5
VLPFC			
L	4.4 ( $\pm$ .9)	5.4 ( $\pm$ 1.1)	8.1 <sup>b</sup>
R	4.2 ( $\pm$ 1.0)	5.2 ( $\pm$ 1.0)	8.9 <sup>b</sup>
DACC			
L	2.7 ( $\pm$ .5)	3.2 ( $\pm$ .9)	4.6 <sup>a</sup>
R	2.8 ( $\pm$ .6)	3.0 ( $\pm$ .7)	.3
VACC			
L	2.0 ( $\pm$ .4)	2.2 ( $\pm$ .7)	1.7
R	2.1 ( $\pm$ .4)	2.4 ( $\pm$ .6)	2.4
Ventral Striatum			
L	1.2 ( $\pm$ .2)	1.1 ( $\pm$ .1)	.8
R	1.2 ( $\pm$ .2)	1.2 ( $\pm$ .2)	.1
Thalamus			
L	6.6 ( $\pm$ .9)	6.5 ( $\pm$ .8)	.3
R	6.8 ( $\pm$ .9)	6.7 ( $\pm$ .9)	.1

Actual ROI/global values are values cited in table  $\times 10^{-3}$ . DLPFC, dorso-lateral prefrontal cortex; L, left; R, right; VLPFC, ventrolateral prefrontal cortex; DACC, dorsal anterior cingulate cortex; VACC, ventral anterior cingulate cortex.

<sup>a</sup> $p < .05$  by analysis of covariance.

<sup>b</sup> $p < .01$  by analysis of covariance.

density, the left ( $Z = 5.38$ ,  $x, y, z = -54, 14, 26$ ,  $p < .0005$ , 300 voxels) and right ( $Z = 6.06$ ,  $x, y, z = 42, 14, 44$ ,  $p < .0005$ , 371 voxels) prefrontal cortices had clusters of voxels with negative main effects of pack-year smoking history (according to the above significance criteria), indicating that greater pack-year smoking history was associated with lower gray matter densities in these regions.


## Discussion

In this study, smokers had smaller relative cortical gray matter volumes and lower gray matter densities than nonsmokers in the prefrontal cortices (DLPFC and VLPFC). Smokers also had smaller left dorsal ACC volumes and lower right cerebellar gray

**Table 2.** Coordinates of Peak Voxels Identified by Voxel-Based Morphometry as Having Lower Gray Matter Densities in Smokers than in Nonsmokers

Region	Z Score	Coordinates			Cluster Size (No. Voxels)
		x	y	z	
Prefrontal Cortex					
Right	6.19	24	42	32	186
	4.75	16	50	32	
	3.49	8	62	28	
	4.46	34	6	48	
	4.36	20	34	52	
Left	3.68	32	18	54	233
	6.66	-32	40	36	
	5.06	-52	30	16	50
Cerebellum					
Right	3.69	16	-56	-26	50

$p < .001$ ,  $Z > 3.39$ , uncorrected; 50 voxel minimum extent.



**Figure 2.** Voxel-based morphometry findings showing brain regions with lower gray matter density in smokers than nonsmokers. Voxels are shown that have significant between-group differences at  $p < .001$ , uncorrected, after applying age and gender as nuisance covariates. DLPFC, dorsolateral prefrontal cortex; VLPFC, ventrolateral prefrontal cortex.

matter densities than nonsmokers. No regions were significantly larger or had significantly higher gray matter densities in smokers compared with nonsmokers. The smoker group also had an association between greater pack-year smoking history and lower prefrontal cortical gray matter density.

Differences between groups in PFC gray matter volume and density might be related to direct effects of cigarettes (Nakamura et al 2000; Stein et al 1998) or cigarette cue exposure (Due et al 2002), predisposing factors that contribute to the development of nicotine dependence (Ernst et al 2001; Mitchell 1999; Spilich et al 1992; Waldeck and Miller 1997), or some combination of these factors. As for the possibility that cigarettes and cigarette cue exposure lead to the group differences found here, nicotine-dependent subjects in this study had between a 9 and 70 pack-year smoking history, meaning that even the lowest pack-year smoking history subject in our study had smoked tens of thousands of cigarettes and been exposed to cigarette-related cues many times before brain scanning. Thus, smoking experience represented a substantial difference between groups, lending support for the theory that smoking (and cigarette cue) exposure might have caused the significant PFC (and other) volume and density differences found here. Animal work supports this theory, with studies of nicotine administration in adolescent rodents resulting in both loss of neurons and changes in nicotinic receptor concentration in the cerebral cortex (Slotkin 2002). The preliminary association found here between greater pack-year smoking history and lower PFC gray matter density strengthens this theory.

As for predisposing factors that might be associated with lower regional brain gray matter in the PFC, prospective and cross-sectional studies of associations between personality traits and smoking indicate that smokers have higher rates than nonsmokers of impulsivity (Mitchell 1999; Waldeck and Miller 1997), neuroticism (Byrne et al 1995; Hu et al 2000), extraversion (Patton et al 1997; Sieber and Angst 1990), antisociality (Barry et al 1997; Patton et al 1997; Reitsma-Street et al 1985), and novelty seeking (Howard et al 1997; Masse and Tremblay 1997). Evidence for these personality traits being associated with smaller PFC volumes is mixed, with behavioral studies indicating PFC function deficits in impulsive individuals (Brower and Price 2001) and MRI studies of antisocial personality indicating smaller prefrontal cortical volumes (Raine et al 2000) but MRI studies of neuroticism (Knutson et al 2001) and impulsivity (Dolan et al

2002) indicating no association of these personality traits with PFC volumes.

The finding of smaller mean left dorsal ACC volumes in smokers might also be related to activation of this region by chronic smoking (Stein et al 1998) or cigarette cue (Brody et al 2002) exposure, or preexisting differences between smokers and nonsmokers. As noted above, smoking leads to changes in behavioral states associated with ACC function, such as improvements in reaction time (Hatsukami et al 1989; Pritchard et al 1992; Shiffman et al 1995), arousal (Parrott and Kaye 1999), motivation (Powell et al 2002), and sustained attention (Rusted et al 2000). Thus, this region might be stimulated repeatedly by smoking or smoking cues, or might be dysfunctional in some individuals (predisposing them to develop nicotine dependence). Furthermore, the fact that the dorsal ACC subregion was significantly smaller in smokers is not surprising. This subdivision is activated during cognitive/motor tasks, such as the Stroop (Whalen et al 1998), whereas smoking history is associated with diminished cognitive flexibility (Kalmijn et al 2002) and functioning (Cervilla et al 2000; Elwood et al 1999; Schinka et al 2002) and reduced psychomotor speed (Kalmijn et al 2002). The cerebellar finding was unexpected, but at least one prior imaging study (Domino et al 2000) did find relative cerebellar activation (in a region overlapping with the one found here) in response to nicotine administration, perhaps indicating chronic stimulation of this region as well.

The findings in this report might aid in the interpretation of studies of brain structure and tissue composition in populations with high rates of nicotine dependence. For example, subjects with depression (Breslau 1995, Breslau et al 1998; de Leon et al 1995, 2002; Glassman et al 1990; Lasser et al 2000), schizophrenia (de Leon et al 1995, 2002; Herran et al 2000; Hughes et al 1986), and other substance dependencies (Bobo 1989; Breslau 1995; Lasser et al 2000) have rates of nicotine dependence much higher than those of the general population. Magnetic resonance imaging studies of these illnesses (Ananth et al 2002; Coffey et al 1993a, 1993b; Fein et al 2002a, 2002b; Hulshoff et al 2002; Kumar et al 1997; Pfefferbaum et al 1998; Suzuki et al 2002) have reported reduced prefrontal cortical gray matter compared with normal control subjects, but they did not focus on the potential confounding role of nicotine dependence. If future prospective studies show that nicotine dependence causes reductions in gray

matter, such findings will point to the need to control for nicotine dependence in studies of these illnesses.

The central limitations of this study were a modest sample size and the absence of sequential MRI scans. A larger, longitudinal study would allow for evaluation of whether volumetric differences between smokers and nonsmokers reflect a biological predisposition to smoking, an effect of smoking, or some combination of these factors. Another limitation is the degree of psychometric characterization of study subjects, which could provide more detail about the complex interplay between smoking, cognitive function, personality characteristics, and regional brain volumes.

Despite these limitations, the present study had several strengths, including the use of relatively stringent thresholds for significance, and similarities in study findings with the use of two independent methods of MRI analysis. Our results indicate significant regional gray matter volume and density differences between smokers and nonsmokers, a potentially important association with smoking behavior that has not yet been widely studied.

*This research was supported by a Veterans Affairs Type I Merit Review Award (ALB), the Tobacco-Related Disease Research Program (ALB [7KT-0098 and 11RT-0024] and EDL [10RT-0091]), and the National Institute on Drug Abuse (ALB [R01 DA15059] and EDL [RO1 DA14093]).*

*We thank Sanjaya Saxena and Richard Traystman for suggestions regarding the manuscript, and Michael Clark for technical assistance in performing magnetic resonance imaging scans. This was presented in part at the American College of Neuropsychopharmacology Annual Meeting, December 9, 2001, Waikoloa, Hawaii, and the European Society for Research on Nicotine and Tobacco Annual Meeting, October 5, 2002, Santander, Spain.*

- Ananth H, Popescu I, Critchley HD, Good CD, Frackowiak RSJ, Dolan RJ (2002): Cortical and subcortical gray matter abnormalities in schizophrenia determined through structural magnetic resonance imaging with optimized volumetric voxel-based morphometry. *Am J Psychiatry* 159:1497–1505.
- Andreasen NC, Flaum M, Swayze V, O'Leary DS, Alliger R, Cohen G, et al (1993): Intelligence and brain structure in normal individuals. *Am J Psychiatry* 150:130–134.
- Ashburner J, Friston K (1997): Multimodal image coregistration and partitioning—a unified framework. *Neuroimage* 6:209–217.
- Ashburner J, Friston KJ (2000): Voxel-based morphometry—the methods. *Neuroimage* 11:805–821.
- Ashburner J, Friston KJ (2001): Why voxel-based morphometry should be used. *Neuroimage* 14:1238–1243.
- Barry KL, Fleming MF, Manwell LB, Copeland LA (1997): Conduct disorder and antisocial personality in adult primary care patients. *J Fam Pract* 45:151–158.
- Bartzokis G, Beckson M, Lu PH, Edwards N, Rapoport R, Wiseman E, et al (2000): Age-related brain volume reductions in amphetamine and cocaine addicts and normal controls: Implications for addiction research. *Psychiatry Res* 98:93–102.
- Bartzokis G, Beckson M, Lu PH, Nuechterlein KH, Edwards N, Mintz J (2001): Age-related changes in frontal and temporal lobe volumes in men—a magnetic resonance imaging study. *Arch Gen Psychiatry* 58:461–465.
- Bench CJ, Frith CD, Grasby PM, Friston KJ, Pauls E, Frackowiak RSJ, et al (1993): Investigations of the functional anatomy of attention using the Stroop test. *Neuropsychologia* 31:907–922.
- Bobo JK (1989): Nicotine dependence and alcoholism epidemiology and treatment. *J Psychoactive Drugs* 21:323–329.
- Bookstein FL (2001): “Voxel-based morphometry” should not be used with imperfectly registered images. *Neuroimage* 14:1454–1462.
- Braver TS, Barch DM, Kelley WM, Buckner RL, Cohen NJ, Miezin FM, et al (2001): Direct comparison of prefrontal cortex regions engaged by working and long-term memory tasks. *Neuroimage* 14:48–59.
- Breslau N (1995): Psychiatric comorbidity of smoking and nicotine dependence. *Behav Genet* 25:95–101.
- Breslau N, Peterson EL, Schultz LR, Chilcoat HD, Andreski P (1998): Major depression and stages of smoking. A longitudinal investigation. *Arch Gen Psychiatry* 55:161–166.
- Brody AL, Mandelkern MA, London ED, Childress AR, Bota RG, Ho ML, et al (2002): Brain metabolic changes during cigarette craving. *Arch Gen Psychiatry* 59:1162–1172.
- Brower MC, Price BH (2001): Neuropsychiatry of frontal lobe dysfunction in violent and criminal behaviour: A critical review. *J Neurol Neurosurg Psychiatry* 71:720–726.
- Burton EJ, Karas G, Paling SM, Barber R, Williams ED, Ballard CG, et al (2002): Patterns of cerebral atrophy in dementia with Lewy bodies using voxel-based morphometry. *Neuroimage* 17:618–630.
- Busatto GE, Garrido GEJ, Almeida OP, Castro CC, Camargo CHP, Cid CG, et al (2003): A voxel-based morphometry study of temporal lobe gray matter reductions in Alzheimer's disease. *Neurobiol Aging* 24:221–231.
- Byrne DG, Byrne AE, Reinhart MI (1995): Personality, stress and the decision to commence cigarette-smoking in adolescence. *J Psychosom Res* 39:53–62.
- Cervilla JA, Prince M, Mann A (2000): Smoking, drinking, and incident cognitive impairment: A cohort community based study included in the Gospel Oak project. *J Neurol Neurosurg Psychiatry* 68:622–626.
- Clarke PBS, Pert C, Pert A (1984): Autoradiographic distribution of nicotine receptors in rat brain. *Brain Res* 323:390–395.
- Coffey CE, Wilkinson WE, Parashos IA, Soady SAR, Sullivan RJ, Patterson LJ, et al (1992): Quantitative cerebral anatomy of the aging human brain: a cross-sectional study using magnetic resonance imaging. *Neurology* 42:527–536.
- Coffey CE, Wilkinson WE, Weiner RD, Parashos IA, Djang WT, Webb MC, et al (1993a): Quantitative cerebral anatomy in depression. A controlled magnetic resonance imaging study. *Arch Gen Psychiatry* 50:7–16.
- Coffey CE, Wilkinson WE, Weiner RD, Ritchie JC, Aque M (1993b): The dexamethasone suppression test and quantitative cerebral anatomy in depression. *Biol Psychiatry* 33:442–449.
- Cowell PE, Turetsky BI, Gur RC, Grossman RI, Shtasel DL, Gur RE (1994): Sex differences in aging of the human frontal and temporal lobes. *J Neurosci* 14:4748–4755.
- Critchley HD, Mathias CJ, Dolan RJ (2001): Neural activity in the human brain relating to uncertainty and arousal during anticipation. *Neuron* 29:537–545.
- Damasio H (1995): *Human Brain Anatomy in Computerized Images*. New York: Oxford University Press.
- Dávila-García MI, Musachio J, Perry D, Xiao Y, Horti A, London E, et al (1997): [<sup>125</sup>I]IPH, an epibatidine analog, binds with high affinity to neuronal nicotinic cholinergic receptors. *J Pharmacol Exp Ther* 282:445–451.
- de Leon J, Dadvand M, Canuso C, White AO, Stanilla JK, Simpson GM (1995): Schizophrenia and smoking: An epidemiologic survey in a state hospital. *Am J Psychiatry* 152:453–455.
- de Leon J, Diaz FJ, Rogers T, Browne D, Dinsmore L (2002): Initiation of daily smoking and nicotine dependence in schizophrenia and mood disorders. *Schizophr Res* 56:47–54.
- Dewey SL, Brodie JD, Gerasimov M, Horan B, Gardner EL, Ashby CRJ (1999): A pharmacologic strategy for the treatment of nicotine addiction. *Synapse* 31:76–86.
- Di Chiara G, Imperato A (1988): Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci U S A* 85:5274–5278.
- Dolan MC, Deakin JF, Roberts N, Anderson IM (2002): Quantitative frontal and temporal structural MRI studies in personality-disordered offenders and control subjects. *Psychiatry Res* 116:133–149.

- Domino EF, Minoshima S, Guthrie S, Ohl L, Ni L, Koeppe RA, et al (2000): Nicotine effects on regional cerebral blood flow in awake, resting tobacco smokers. *Synapse* 38:313–321.
- Due DL, Huettel SA, Hall WG, Rubin DC (2002): Activation in mesolimbic and visuospatial neural circuits elicited by smoking cues: Evidence from functional magnetic resonance imaging. *Am J Psychiatry* 159:954–960.
- Elwood PC, Gallacher JEJ, Hopkinson CA, Pickering J, Rabbitt P, Stollery B, et al (1999): Smoking, drinking, and other life style factors and cognitive function in men in the Caerphilly cohort. *J Epidemiol Community Health* 53:9–14.
- Ernst M, Heishman SJ, Spurgeon L, London ED (2001): Smoking history and nicotine effects on cognitive performance. *Neuropsychopharmacology* 25:313–319.
- Fagerström KO (1978): Measuring the degree of physical dependence to tobacco smoking with reference to individualization of treatment. *Addict Behav* 3:235–241.
- Fein G, Di Sclafani V, Cardenas VA, Goldmann H, Tolou-Shams M, Meyerhoff DJ (2002a): Cortical gray matter loss in treatment-naive alcohol dependent individuals. *Alcohol Clin Exp Res* 26:558–564.
- Fein G, Di Sclafani V, Meyerhoff DJ (2002b): Prefrontal cortical volume reduction associated with frontal cortex function deficit in 6-week abstinent crack-cocaine dependent men. *Drug Alcohol Depend* 68:87–93.
- First MB, Spitzer RL, Gibbon M, Williams JBW (1995): *Structured Clinical Interview for DSM-IV Axis I Disorders—Patient Edition, version 2.0*. Washington, DC: American Psychiatric Press.
- Franklin TR, Acton PD, Maldjian JA, Gray JD, Croft JR, Dackis CA, et al (2002): Decreased gray matter concentration in the insular, orbitofrontal, cingulate, and temporal cortices of cocaine patients. *Biol Psychiatry* 51:134–142.
- Friston KJ, Worsley K, Frackowiak RSJ, Mazziotta J, Evans AC (1994): Assessing the significance of focal activations using their spatial extent. *Hum Brain Mapp* 1:214–220.
- Fukuda H, Kitani M (1996): Cigarette smoking is correlated with the periventricular hyperintensity grade on brain magnetic resonance imaging. *Stroke* 27:645–649.
- Glassman AH, Helzer JE, Covey LS, Cottler LB, Stetner F, Tipp JE, et al (1990): Smoking, smoking cessation, and major depression. *JAMA* 264:1546–1549.
- Good CD, Scahill RI, Fox NC, Ashburner J, Friston KJ, Chan D, et al (2002): Automatic differentiation of anatomical patterns in the human brain: Validation with studies of degenerative dementias. *Neuroimage* 17:29–46.
- Gur RC, Gunning-Dixon F, Bilker WB, Gur RE (2002a): Sex differences in temporo-limbic and frontal brain volumes of healthy adults. *Cereb Cortex* 12:998–1003.
- Gur RC, Gunning-Dixon FM, Turetsky BI, Bilker WB, Gur RE (2002b): Brain region and sex differences in age association with brain volume: A quantitative MRI study of healthy young adults. *Am J Geriatr Psychiatry* 10:72–80.
- Hamilton M (1967): Development of a rating scale for primary depressive illness. *Br J Soc Psychol* 6:278–296.
- Hamilton M (1969): Diagnosis and rating of anxiety. *Br J Psychiatry* 3:76–79.
- Hatsukami D, Fletcher L, Morgan S, Keenan R, Amble P (1989): The effects of varying cigarette deprivation duration on cognitive and performance tasks. *J Subst Abuse* 1:407–416.
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO (1991): The Fagerström Test for Nicotine Dependence: A revision of the Fagerström Tolerance Questionnaire. *Br J Addict* 86:1119–1127.
- Herran A, de Santiago A, Sandoya M, Fernandez MJ, Diez-Manrique JF, Vazquez-Barquero JL (2000): Determinants of smoking behaviour in outpatients with schizophrenia. *Schizophr Res* 41:373–381.
- Horti AG, Scheffel U, Kimes A, Musachio J, Ravert H, Mathews W, et al (1998): Synthesis and evaluation of N-<sup>11</sup>C]methylated analogues of epibatidine as tracers for positron emission tomographic studies of nicotinic acetylcholine receptors. *J Med Chem* 41:4199–4206.
- Howard G, Wagenknecht LE, Cai JW, Cooper L, Kraut MA, Toole JF (1998): Cigarette smoking and other risk factors for silent cerebral infarction in the general population. *Stroke* 29:913–917.
- Howard MO, Kivlahan D, Walker RD (1997): Cloninger's tridimensional theory of personality and psychopathology: Applications to substance use disorders. *J Stud Alcohol* 58:48–66.
- Hu S, Brody CL, Fisher C, Gunzerath L, Nelson ML, Sabol SZ, et al (2000): Interaction between the serotonin transporter gene and neuroticism in cigarette smoking behavior. *Mol Psychiatry* 5:181–188.
- Hughes JR, Hatsukami DK, Mitchell JE, Dahlgren LA (1986): Prevalence of smoking among psychiatric outpatients. *Am J Psychiatry* 143:993–997.
- Hulshoff HE, Schnack HG, Bertens MGBC, van Haren NEM, van der Tweel I, Staal WG, et al (2002): Volume changes in gray matter in patients with schizophrenia. *Am J Psychiatry* 159:244–250.
- Jacobsen LK, Giedd JN, Gottschalk C, Kosten TR, Krystal JH (2001a): Quantitative morphology of the caudate and putamen in patients with cocaine dependence. *Am J Psychiatry* 158:486–489.
- Jacobsen LK, Giedd JN, Kreek MJ, Gottschalk C, Kosten TR (2001b): Quantitative medial temporal lobe brain morphology and hypothalamic-pituitary-adrenal axis function in cocaine dependence: A preliminary report. *Drug Alcohol Depend* 62:49–56.
- Jernigan TL, Archibald SL, Berhow MT, Sowell ER, Foster DS, Hesselink JR (1991): Cerebral structure on MRI, part I: Localization of age-related changes. *Biol Psychiatry* 29:55–67.
- Jernigan TL, Archibald SL, Fennema-Notestine C, Gamst AC, Stout JC, Bonner J, et al (2001): Effects of age on tissues and regions of the cerebrum and cerebellum. *Neurobiol Aging* 22:581–594.
- Kalmijn S, van Boxtel MPJ, Verschuren MWM, Jolles J, Launer LJ (2002): Cigarette smoking and alcohol consumption in relation to cognitive performance in middle age. *Am J Epidemiol* 156:936–944.
- Karas G, Rombouts SARB, van Schijndel RA, Powels P, Scheltens P, Barkhof F, et al (2002): Patterns of cerebral atrophy detected by voxel-based morphometry (VBM) in Alzheimer's disease. *Neurobiol Aging* 23:S352–S353.
- Knutson B, Momenan R, Rawlings RR, Fong GW, Hommer D (2001): Negative association of neuroticism with brain volume ratio in healthy humans. *Biol Psychiatry* 50:685–690.
- Kobayashi S, Okada K, Koide H, Bokura H, Yamaguchi S (1997): Subcortical silent brain infarction as a risk factor for clinical stroke. *Stroke* 28:1932–1939.
- Kumar A, Schweizer E, Jin ZS, Miller D, Bilker W, Sean LL, et al (1997): Neuro-anatomical substrates of late-life minor depression: A quantitative magnetic resonance imaging study. *Arch Neurology* 54:613–617.
- Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH (2000): Smoking and mental illness: A population-based prevalence study. *JAMA* 284:2606–2610.
- Liao DP, Cooper L, Cai JW, Toole J, Bryan N, Burke G, et al (1997): The prevalence and severity of white matter lesions, their relationship with age, ethnicity, gender, and cardiovascular disease risk factors: The ARIC study. *Neuroepidemiology* 16:149–162.
- Liu X, Matochik JA, Cadet JL, London ED (1998): Smaller volume of prefrontal lobe in polysubstance abusers: A magnetic resonance imaging study. *Neuropsychopharmacology* 18:243–252.
- London ED, Scheffel U, Kimes AS, Kellar KJ (1995): In vivo labeling of nicotinic acetylcholine receptors in brain with [<sup>3</sup>H]epibatidine. *Eur J Pharmacol* 278:R1–R2.
- London ED, Waller SB, Wamsley JK (1985): Autoradiographic localization of [<sup>3</sup>H]nicotine binding sites in the rat brain. *Neurosci Lett* 53:179–184.
- Longstreth WT, Arnold AM, Manolio TA, Burke GL, Bryan N, Jungreis CA, et al (2000): Clinical correlates of ventricular and sulcal size on cranial magnetic resonance imaging of 3,301 elderly people: The cardiovascular health study. *Neuroepidemiology* 19:30–42.
- Longstreth WT, Bernick C, Manolio TA, Bryan N, Jungreis CA, Price TR (1998): Lacunar infarcts defined by magnetic resonance imaging of 3660 elderly people: The cardiovascular health study. *Arch Neurology* 55:1217–1225.
- Longstreth WT, Diehr P, Manolio TA, Beauchamp NJ, Jungreis CA, Lefkowitz D (2001): Cluster analysis and patterns of findings on cranial magnetic resonance imaging of the elderly: The cardiovascular health study. *Arch Neurology* 58:635–640.
- Mai JK, Assheuer J, Paxton DA (1997): *Atlas of the Human Brain*. San Diego: Academic Press, Harcourt Brace and Company.
- Masse LC, Tremblay RE (1997): Behavior of boys in kindergarten and the onset of substance use during adolescence. *Arch Gen Psychiatry* 54:62–68.
- Meyer-Lindenberg A, Poline JB, Kohn PD, Holt JL, Egan MF, Weinberger DR, et al (2001): Evidence for abnormal cortical functional connectivity during working memory in schizophrenia. *Am J Psychiatry* 158:1809–1817.
- Mitchell SH (1999): Measures of impulsivity in cigarette smokers and non-smokers. *Psychopharmacology* 146:455–464.
- Mueller EA, Moore MM, Kerr DC, Sexton G, Camicioli RM, Howieson DB, et al (1998): Brain volume preserved in healthy elderly through the eleventh decade. *Neurology* 51:1555–1562.

- Murphy DG, DeCarli C, McIntosh AR, Daly E, Mentis MJ, Pietrini P, et al (1996): Sex differences in human brain morphometry and metabolism: An in vivo quantitative magnetic resonance imaging and positron emission tomography study on the effect of aging. *Arch Gen Psychiatry* 53:585–594.
- Musachio J, Villemagne V, Scheffel U, Stathis M, Finley P, Horti A, et al (1997): [<sup>125</sup>I/<sup>123</sup>I] IPH: A radioiodinated analog of epibatidine for in vivo studies of nicotinic acetylcholine receptors. *Synapse* 26:392–399.
- Nakamura H, Tanaka A, Nomoto Y, Ueno Y, Nakayama Y (2000): Activation of fronto-limbic system in the human brain by cigarette smoking: Evaluated by a CBF measurement. *Keio J Med* 49(suppl 1):A122–A124.
- Oldfield RC (1971): The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* 9:97–113.
- Parrott AC, Kaye FJ (1999): Daily uplifts, hassles, stresses and cognitive failures: In cigarette smokers, abstaining smokers, and non-smokers. *Behav Pharmacol* 10:639–646.
- Patton D, Barnes GE, Murray RP (1997): A personality typology of smokers. *Addict Behav* 22:269–273.
- Pfefferbaum A, Sullivan EV, Rosenbloom MJ, Mathalon H, Lim KO (1998): A controlled study of cortical gray matter and ventricular changes in alcoholic men over a 5-year interval. *Arch Gen Psychiatry* 55:905–912.
- Powell J, Dawkins L, Davis RE (2002): Smoking, reward responsiveness, and response inhibition: Tests of an incentive motivational model. *Biol Psychiatry* 51:151–163.
- Pritchard WS, Robinson JH, Guy TD (1992): Enhancement of continuous performance task reaction-time by smoking in nondeprived smokers. *Psychopharmacology* 108:437–442.
- Raine A, Lencz T, Bihle S, LaCasse L, Colletti P (2000): Reduced prefrontal gray matter volume and reduced autonomic activity in antisocial personality disorder. *Arch Gen Psychiatry* 57:119–127.
- Rauch SL, Shin LM, Dougherty DD, Alpert NM, Orr SP, Lasko M, et al (1999): Neural activation during sexual and competitive arousal in healthy men. *Psychiatry Res* 91:1–10.
- Raz N, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, et al (1997): Selective aging of the human cerebral cortex observed in vivo: Differential vulnerability of the prefrontal gray matter. *Cereb Cortex* 7:268–282.
- Reiss AL, Abrams MT, Singer HS, Ross JL, Denckla MB (1996): Brain development, gender and IQ in children: A volumetric imaging study. *Brain* 119:1763–1774.
- Reitsma-Street M, Offord DR, Finch T (1985): Pairs of same-sexed siblings discordant for antisocial behavior. *Br J Psychiatry* 146:415–423.
- Rose JE, Behm FM, Westman EC, Mathew RJ, London ED, Hawk TC, et al (2003): PET studies of the influences of nicotine on neural systems in cigarette smokers. *Am J Psychiatry* 160:323–333.
- Rusted JM, Caulfield D, King L, Goode A (2000): Moving out of the laboratory: Does nicotine improve everyday attention? *Behav Pharmacol* 11:621–629.
- Salat DH, Kaye JA, Janowsky JS (2001): Selective preservation and degeneration within the prefrontal cortex in aging and Alzheimer disease. *Arch Neurol* 58:1403–1408.
- Schinka JA, Vanderploeg RD, Rogish M, Ordorica PI (2002): Effects of alcohol and cigarette use on cognition in middle-aged adults. *J Int Neuropsychol Soc* 8:683–690.
- Shiffman S, Paty JA, Gnys M, Elash C, Kassel JD (1995): Nicotine withdrawal in chippers and regular smokers: Subjective and cognitive effects. *Health Psychol* 14:301–309.
- Sieber MF, Angst J (1990): Alcohol, tobacco and cannabis: 12-year longitudinal associations with antecedent social-context and personality. *Drug Alcohol Depend* 25:281–292.
- Slotkin TA (2002): Nicotine and the adolescent brain: Insights from an animal model. *Neurotoxicol Teratol* 24:369–384.
- Spilich GJ, June L, Renner J (1992): Cigarette-smoking and cognitive performance. *Br J Addict* 87:1313–1326.
- Stein E, Pankiewicz J, Harsch HH, Cho JK, Fuller SA, Hoffmann RG, et al (1998): Nicotine-induced limbic cortical activation in the human brain: A functional MRI study. *Am J Psychiatry* 155:1009–1015.
- Steinmetz H, Seitz RJ (1991): Functional anatomy of language processing: Neuroimaging and the problem of individual variability. *Neuropsychologia* 29:1149–1161.
- Suzuki M, Nohara S, Hagino H, Kurokawa K, Yotsutsuji T, Kawasaki Y, et al (2002): Regional changes in brain gray and white matter in patients with schizophrenia demonstrated with voxel-based analysis of MRI. *Schizophrenia Res* 55:41–54.
- Tisserand D, Pruessner J, Sanz Arigita E, van Boxtel M, Evans A, Jolles J, et al (2002): Regional frontal cortical volumes decrease differentially in aging: An MRI study to compare volumetric approaches and voxel-based morphometry. *Neuroimage* 17:657–659.
- Tisserand DJ, Visser PJ, van Boxtel MP, Jolles J (2000): The relation between global and limbic brain volumes on MRI and cognitive performance in healthy individuals across the age range. *Neurobiol Aging* 21:569–576.
- Tsushima Y, Tanizaki Y, Aoki J, Endo K (2002): MR detection of microhemorrhages in neurologically healthy adults. *Neuroradiology* 44:31–36.
- Van Der Werf YD, Tisserand DJ, Visser PJ, Hofman PA, Vuurman E, Uylings HB, et al (2001): Thalamic volume predicts performance on tests of cognitive speed and decreases in healthy aging. A magnetic resonance imaging-based volumetric analysis. *Brain Res Cogn Brain Res* 11:377–385.
- Waldeck TL, Miller LS (1997): Gender and impulsivity differences in licit substance use. *J Subst Abuse* 9:269–275.
- Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, Jenike MA, et al (1998): The emotional counting Stroop paradigm: A functional magnetic resonance imaging probe of the anterior cingulate affective division. *Biol Psychiatry* 44:1219–1228.
- Yamashita K, Kobayashi S, Yamaguchi S, Koide H (1996): Cigarette smoking and silent brain infarction in normal adults. *Intern Med* 35:704–706.
- Yetkin FZ, Fischer ME, Papke RA, Haughton VM (1993): Focal hyperintensities in cerebral white matter on MR images of asymptomatic volunteers: Correlation with social and medical histories. *AJR Am J Roentgenol* 161:855–858.
- Zubieta J, Lombardi U, Minoshima S, Guthrie S, Ni L, Ohl LE, et al (2001): Regional cerebral blood flow effects of nicotine in overnight abstinent smokers. *Biol Psychiatry* 49:906–913.
- Zurowski B, Gostomzyk J, Gron G, Weller R, Schirrmeyer H, Neumeier B, et al (2002): Dissociating a common working memory network from different neural substrates of phonological and spatial stimulus processing. *Neuroimage* 15:45–57.